



## Sedation with medicinal plants: A review of medicinal plants with sedative properties in Iranian ethnobotanical documents

Fariba Bahmani<sup>1</sup>, Hamidreza Kazemeini<sup>2</sup>, Fatemeh Hoseinzadeh-Chahkandak<sup>3</sup>, Tahereh Farkhondeh<sup>4\*</sup>, Mahshid Sedaghat<sup>5</sup>

<sup>1</sup>Shohada Hospital, Dehloran City, Ilam University of Medical Sciences, Ilam, Iran

<sup>2</sup> Assistant Professor, Department of Food Hygiene, Faculty of Veterinary Medicine, Amol University of Special Modern Technologies, Amol, Iran

<sup>3</sup>Social Determinants of Health Research Center, Birjand University of Medical Sciences, Birjand, Iran

<sup>4</sup> Cardiovascular Diseases Research Center, Birjand University of Medical Sciences, Birjand, Iran

<sup>5</sup>Faculty of Medicine, Birjand University of Medical Sciences, Birjand, Iran

### \*Correspondence to

Dr. Tahereh Farkhondeh  
farkhondeh2324@gmail.com

**Received:** 11 August 2019

**Accepted:** 18 October 2019

**ePublished:** 20 November 2019

**Keywords:** Sedative, Medicinal plants, Ethnobotany, Iran.

**Citation:** Bahmani F, Kazemeini HR, Hoseinzadeh-Chahkandak F, Farkhondeh T, Sedaghat M. Sedation with medicinal plants: A review of medicinal plants with sedative properties in Iranian ethnobotanical documents. *Plant Biotechnol Persa* 2019; 1(1): 13-24.

### Abstract

The complexity of today's modern society, increasing problems and changes in lifestyles can lead to stress and anxiety in people. To control stress and anxiety, sedative drugs that have anxiolytic and sedative effects and temporarily calm the person and lead to a pleasant sleep are used. However, it must also be kept in mind that each of these drugs have their own side effects. The information in this review study was obtained from scientific publications indexed in databases such as the Information Sciences Institute, PubMed, Scopus, Scientific Information Database, Magiran, and Google Scholar that were retrieved by using relevant search terms including sedative, ethnobotany, herbal drug, Iran, medicinal plants, extracts, and essential oils. A total of 46 articles were retrieved by our initial search. The articles were reviewed. Forty three articles were further reviewed, 16 of which were finally used to review the sedative effects in ethnobotany. Being sedative is one of the reported properties of some herbal drugs. The use of these materials can be beneficial and will not cause side effects like those of conventional sedative drugs. The use of oral herbal drugs is one of the ancient methods that have been used to control anxiety, stress and pain. Besides, numerous side effects of chemical drugs have caused a renewed tendency to using herbal medicine. Therefore, this study is aimed to introduce some of the most effective medicinal plants with sedative properties based on Iranian ethnobotanical documents.

## Introduction

The complexity of today's modern society, increasing problems, and changes in lifestyles can lead to stress, anxiety and mental disorders in people that have become the most important problems in modern societies [1-5]. Stress is defined according to mental perceptions of an individual and his/her perception of a situation as being beyond his/her abilities that can disturb his/her health. In general, stress is experienced by all people and it should be noted that certain degrees of the stress can be effective in increasing and improving the performance of individuals, but when stress exceeds the capacity of a person, behavioral and emotional disorders occur in him/her [6,7]. Anxiety is a type of natural feeling that is experienced in threatening situations. Clinical properties of anxiety disorders are the same as normal anxiety and do not differ in type, but do in duration and severity [8]. Mental state, anxiety and concern are accompanied by certain physiological changes such as increased heart rate, hypertension, increased respiratory rate, and increased muscle tone, making the individual ready for escape or defense [9]. Pathologic anxiety is one of the most common psychiatric disorders that cause disturbance of daily life and suffering in humans. Benzodiazepines and barbiturates are used to treat this condition [10]. Sedatives are drugs that, by suppressing the central nervous system (CNS), can lead to calmness, relaxation of the nervous system, relief of anxiety and stress, sleepiness, euphoria, decrease in respiratory rate, sensations of imbalance, muscle relaxation, impaired

judgments, and slow and hesitant reactions [11,12]. Although taking chemical medications to relieve anxiety can temporarily calm the person and bring about a pleasant sleep for him/her, it must also be kept in mind that each of these medications have their own side effects, and it is therefore advised to take them according to the prescription of a physician and not to take them at high doses under any circumstances [11]. Sedatives have anxiolytic and calming effects and are also used to treat insomnia. Common sedative medications include benzodiazepines, opiates, barbiturates and hypnotics. Opiates are one of the most sought-after drug classes because of causing analgesia. In addition to analgesia, these drugs also have sedative effects [13,14]. Side effects of sedatives, especially in the elderly, are due to hypersensitivity to these drugs, which can lead to certain complications such as weakened response of the nervous system to hypercapnia, hypoxia, hypotension that should be treated, and postoperative apnea [15-18]. Medicinal plants and foods have a huge impact on the health of individuals [19,20]. Any food has a specific impact on the function of the body. *Being sedative* is one of the reported properties of some herbal drugs. They are very beneficial and less likely to cause the side effects of conventional sedative drugs. The use of oral herbal drugs is one of the ancient methods that have been used to control anxiety, stress and pain. In addition, several side effects due to chemical drugs have caused a renewed tendency to using herbal medicine [21]. Medicinal plants have been shown to boost the immune system, produce antimicrobial effects,

and be effective on infectious and non-infectious diseases and chronic diseases, which have been proven in both ethnobotanical and pharmacological studies [22-27]. Therefore, this study is aimed to introduce some of the most effective medicinal plants with sedative properties based on Iranian ethnobotanical documents.

### Methodology

The information in this review study was obtained from scientific publications indexed in databases such as the Information Sciences Institute, PubMed, Scopus, Scientific Information Database, Magiran, and Google Scholar that were retrieved by using relevant search terms including sedative, ethnobotany, herbal drug, Iran, medicinal plants, extracts, and essential oils. A total of 46 articles were retrieved by our initial search. The articles were reviewed. Forty three articles were further reviewed, 16 of which were finally used to review the sedative effects in ethnobotany.

### Results

Ethnobotany is the study of how people of a particular ethnic group, culture, or region use indigenous plants. The results of this review of Iranian ethnobotanical documents showed that medicinal plants from certain families such as Apiaceae, Asteraceae and Lamiaceae are among the most important plants used in Iran for their sedative effects (Table 1).

**Table 1. Medicinal plants with sedative effects according to Iranian ethnobotanical documents**

Scientific name	Family name	Domestic name	Used organs	Region
<i>Allium sp</i>	Liliaceae	Paiz vahshi	Bulb	Abadeh fars [28]
<i>Nonnea cospica</i>	Boraginaceae	Chesh gorbei	Flower	Joupar kerman [29]
<i>Valeriana ficariaefolia</i>	Valerianaceae	Alafe gorbeh	Root, rizhome	Joupar kerman [29]
<i>Avena sativa</i>	Poaceae	Jo dosar	Aerial parts	Zabol [30]
<i>Rubia tinctorum</i>	Rubiaceae	Ronas	Root	Zabol [30]
<i>Heracleum persicum</i>	Apiaceae	Golpar	Fruit and leaf	Khuzistan [31]

<i>Kelussia odoratissima</i>	Apiaceae	Karafseh kouhi	Aerial parts	Khuzistan [31]
<i>Pimpinella anisum</i>	Apiaceae	Anison	Fruit	Khuzistan [31]
<i>Trachyspermum copticum</i>	Apiaceae	Zenian	Fruit and seed	Khuzistan [31]
<i>Centurea depressa</i>	Asteraceae	Gole gandom	Stem	Khuzistan [31]
<i>Lactuca virosa</i>	Asteraceae	Kahoye vahshi	Aerial parts	Khuzistan [31]
<i>Echium amoenum</i>	Boraginaceae	Gav zaban	Flower	Khuzistan [31]
<i>Brassica napus</i>	Brassicaceae	Shalgham	Seed and root	Khuzistan [31]
<i>Humulus lupulus</i>	Cannabaceae	Razak	Inflorescence	Khuzistan [31]
<i>Lavandula angustifolia</i>	Lamiaceae	Ostokhodous	Flower, stem and leaf	Khuzistan [31]
<i>Melissa officinalis</i>	Lamiaceae	Badranjbouyeh	Leaf and stem	Khuzistan [31]
<i>Mentha longifolia</i>	Lamiaceae	Pouneh	Leaf and stem and flower	Khuzistan [31]
<i>Scutellaria lateviflora</i>	Lamiaceae	Varangbou	Seed	Khuzistan [31]
<i>Peganumharmala</i>	Nitrariaceae	Spand	Seed and fruit	Khuzistan [31]
<i>Crataegus curvisepala</i>	Rosaceae	Zalzalak	Leaves, flowers and fruits	Khuzistan [31]
<i>Hyoscyamus orthocarpus</i>	Solanaceae	Bazrolbanj	Leaf and seed	Khuzistan [31]
<i>Pistacia khinjuk</i>	Anacardiaceae	Pesteh kouhi	Fruit	Fasa [32]
<i>Avenasativa</i>	Graminaceae	Jodosaar	Fruit	Fasa [32]
<i>Stachysinflata</i>	Lamiaceae	Sonbole badkonaki	Flowered flower	Fasa [32]
<i>Alliumcanadense</i>	Liliaceae	Sirmouk	Aerial parts	Fasa [32]
<i>Papaverdubium</i>	Papaveraceae	Shaghayegh	Flower & Gum	Fasa [32]
<i>Plantagomajor</i>	Plantaginaceae	Barhang	Seed and leaf	Fasa [32]
<i>Citrusaurantium</i>	Rutaceae	Naranj	Flower and fruit	Fasa [32]

<i>Marrubium vulgare</i>	Labiatae	Ferasion	Stem and leaf	Gichak [32]
<i>Zataria multiflora</i>	Lamiaceae	Abeshm	Aerial parts	Naein [33]
<i>Ziziphora tenuior</i>	Lamiaceae	Ostaghodos	Aerial parts	Naein [34]
<i>Echium spp.</i>	Boraginaceae	Gavzaban	Flower	Erim neka [35]
<i>Viola ororata</i>	Violaceae	Banafsheh	Flower	Erim neka [35]
<i>Achillea biebersteinii</i>	Asteraceae	Boumadaran	Leaf and flower	Dehloran and Abdanan [36]
<i>Cerasus microcarpa</i>	Rosaceae	Albalouyeh vahshi	Resin	Dehloran and Abdanan [36]
<i>Hypericum scabrum</i>	Hypericaceae	Gole raei	Flowered flower	Dehloran and Abdanan [36]
<i>Lonicera nummulariifolia</i>	Caprifoliaceae	Pilakhor	Leaf and flower	Dehloran and Abdanan [36]
<i>Narcissus tazetta</i>	Amaryllidaceae	Narges	Flower	Dehloran and Abdanan [36]
<i>Papaver dubium</i>	Papaveraceae	Khashkhash tanaz	Leaf and flower	Dehloran and Abdanan [36]
<i>Stachys lavandulifolia</i>	Lamiaceae	Sonbolei	Leaf and flower	Dehloran and Abdanan [36]
<i>Grammosciadium platycarpum</i>	Apiaceae	Shevid kouhi	Aerial parts	Behbahan [37]
<i>Anthemis cotula</i>	Apiaceae	Babouneh bahari	Aerial parts	Behbahan [37]
<i>Leonurus cardiaca</i>	Lamiaceae	Ostoghodos vahshi	Aerial parts	Behbahan [37]
<i>Mentha mozaffarianii</i>	Lamiaceae	Pouneh kouhi	Aerial parts	Behbahan [37]
<i>Datura stramonium</i>	Solanaceae	Tatoureh	Aerial parts	Behbahan [37]
<i>Celtis australis</i>	Ulmaceae	Daghdaghan	Aerial parts	Behbahan [37]

<i>Vitex pseudonegundo</i>	Verbenaceae	Panj angosht	Aerial parts	Behbahan [37]
<i>Verbenaceae</i>	Verbenaceae	Gole gavzaban	Aerial parts	Razovergolan [38]
<i>Cannabis sativa</i>	Cannabaceae	Shahdaneh	Leaf, seed	Sirjan [39]
<i>Echium amoenum</i>	Boraginaceae	Zabangav	Aerial parts	Sirjan [39]
<i>Datura innoxia</i>	Solanaceae	Datoureh	Stem	Sistan [40]
<i>Solanumnigrum</i>	Solanaceae	Tajrizi siah	Leaf and seed	Sistan [40]
<i>Borago officinalis</i>	Boraginaceae	Sisenak	Flower and leaf	North Iran [41]
<i>Hyoscyamus niger</i>	Solanaceae	Taltouleh	Leaf	North Iran [41]
<i>Melissa officinalis</i>	Lamiaceae	Badranjbouyeh	Aerial parts	Kazeroun [42]
<i>Vitex agnus-castus</i>	Verbenaceae	Bangela	Leaf and fruit	Hormozgam [43]

## Discussion

Depression, fatigue, anxiety, anger, annoyance, chastity and agitation These familiar traits nowadays are heard from every man and woman, old and young, and even children and adolescents. Relaxation or mild anesthesia is a method that reduces the amount of pain and incitement caused by the therapeutic process or painful diagnosis using sedative medications. Drugs that are used in this process include Propofol, Probiotics, Ketamine, Fentanyl, and Midazolam [44]. Over the years, natural medicines, especially medicinal herbs, have been the basis and sometimes even the only remedy, while raw materials used in the pharmaceutical industry were used [45]. Today, due to the adverse effects of some chemical drugs and people's beliefs about the increased safety of the use of natural compounds, the tendency to use herbal medicines has increased. One of the

problems that is spreading among our society today is insomnia and neurotic stress caused by personal and social problems. This has led people to tend to use sedative medications.

Among the others, the most effective plants with sedative hypnotic properties are *Valeriana officinalis*. It is noted as a remedy to modulate the troubled nerves and to induce untroubled sleep. It has mild effects, however, it is a safe remedy for use [46]. *Valeriana officinalis* is prepared as ethanol or aqueous extracts. Its therapeutic dose is about 600 mg per day. The mechanism action of *V. officinalis* has been roughly identified previously. However, the plant has more than 100 components and it is not known which one is responsible for its anxiolytic action [47]. Valeric and iso-valeric acids are among the most important components and have GABA re-uptake inhibitory in brain, which are involved in its

tranquilizer effects. Valerian preparations of usually do not impose immediate action and usually 2 to 4 weeks is necessary for achievement significant improvements [48]. The main value of this plant is its capability of promoting natural sleep. However, this will be achieved after several weeks consumption of this remedy. It has no adverse health effects or risk of dependence [49]. Echim amoenum is another plant which possess sedative effects. Chemical substances such as flavonoids, saponins and unsaturated terpenoids and sterols has been identified in Echim amoenum. Natural Flavonoids and their derivatives maintain mild sedative and anxiolytic effects via binding to the central benzodiazepine receptors [50]. Infusion of hydroalcoholic and aqueous extracts of Echim amoenum in mice have been reported to induce anxiolytic activities [51]. Furthermore, it has been showed that administration of 50 mg/kg of Echim amoenum extract induces anxiolytic effects and a reduction in locomotor activity [52]. Another noteworthy plant to mention is humulus lupus. It is used in conditions such as insomnia and anxiety. Humulus lupus extracts contain alpha bitter acids and beta acids which both have sedative effects, but the latter is weaker compared to the first one. Humulus lupus interacts with melatonin receptors which play a role in sleep [53]. In a study by shishehgar et al. 2012, it was showed that humulus lupulus had more sedation, pre-anesthetic, and anti-anxiety effect compared to diazepam [54]. It should be noted that high level of free radicals, which usually lead to oxidative stress and neurodegenerative disorders and health

complaints. Patients with primary insomnia have been shown to possess significantly lower glutathione activity and higher amount of blood malondialdehyde than normal subjects which show that insomnia increases oxidative stress and sleep can reduce oxidative stress [55]. The plants presented in this review and a lot of other plants have antioxidant activities [56-58]. Therefore, these plants may also show positive effects on insomnia. It is important to note that most of plants have multi-function [59-61].

### **Conclusion**

Therefore, these plants may also have beneficial effect on people with other problems, too.

### **Funding**

The authors received no funding

### **Conflict of interest**

The authors declared that they have no conflicts of interest.

### **Ethics approval**

Not required

### **Funding/Support**

None.

### **References**

- [1] Bhui K, Fletcher A. Common mood and anxiety states: gender differences in the protective effect of physical activity. *Soc Psychiatry Psychiatr Epidemiol.* 2000;35(1):28–35.

- [2] Valizadeh R, Veisani Y, Delpisheh A, Kikhavani S, Sohrabnejad A. Major depression and psychiatric disorders in Iranian prisoners based on a clinical interview: A systematic review and meta-analysis. *Shiraz E Med J.* 2017; 18(6):e44979.
- [3] Mohammadalizadeh Charandabi S, Rezaei N, Hakimi S, Khatami SH, valizadeh R, Azadi A. Sleep disturbances and sexual function among men aged 45-75 years in an urban area of Iran. *Sleep Sci.* 2016; 9(1): 29–34.
- [4] Valizadeh R, Sarokhani D, Sarokhani M, Sayehmiri K, Ostovar R, Angh P, Malekzadeh M. A study of prevalence of anxiety in Iran: Systematic review and meta-analysis. *Der Pharma Chemica* 2016; 8(21): 48-57.
- [5] Motl RW, Birnbaum AS, Kubik MY, Dishman RK. Naturally occurring changes in physical activity are inversely related to depressive symptoms during early adolescence. *Psychosom Med.* 2004; 66: 336-342.
- [6] Pourghane P, Sharif azar E, Zaer sabet F and Khorsandi M. Survey the effect of religious beliefs in stress reduction in students of Langroud Faculty of Medical Sciences. *Guilan Uni Med Sci J.* 2011; 20 (63): 10-5.
- [7] Hakimgavadi M, Lavasani M, Haghightgoo M and Zebardast O. Relation of depression, anxiety, stress and personality trait in dedicated, s children. *Veteran Med J.* 2011; 3 (9): 9 – 16.
- [8] KE. WHSAM. Antioxidant activity in extracts from coriander. *Food Chem.* 2004; 88 (2): 293-7.
- [9] Linden C. Anxiety and Panic Attacks Solution: Reprogramming your anxiety with the Linden Method. Available from: <http://www.Lindenmethod.com>
- [10] Finn DA, Rutledge-Gorman MT, Crabbe JC. Genetic animal models of anxiety. *Neurogenetics.* 2003; 4(3):109-35.
- [11] Sadock BJ, Sadock VA. Kaplan & Sadock's Comprehensive text book of Psychiatry. Seventh Edition, Lippincott Williams & Wilkins Co, Volume 1, 2005; 1720-1729.
- [12] Guyton AC, Hall JE. Textbook of Medical Physiology, 11st ed. Philadelphia; Elsevier Saunders Co. 2006; 739-46.
- [13] Ibrahim EH, Kollef MH. Using protocols to improve the outcomes of mechanically ventilated patients. Focus on weaning and sedation. *Crit Care Clin.* 2001; 17(4): 989-1001.
- [14] Shapiro BA, Peruzzi WT. Respiratory care. In: Miller RD, ed: Anesthesia, 5th edition, New York: Churchill Livingstone; 2000. 4: 2424.
- [15] Rivera, R. and J.F. Antognini, Perioperative drug therapy in elderly patients. *J Am Soc Anesthesiol.* 2009; 110(5):1176-81.
- [16] Lawrence VA, Cornell JE, Smetana GW. Strategies to Reduce Postoperative Pulmonary Complications after Noncardiothoracic Surgery: Systematic Review for the American College of Physicians Strategies to Reduce Postoperative



Pulmonary Complications after Noncardiothoracic Surgery. *Annals Int Med.* 2006. 144(8): p. 596-608.

[17] Hasukić Š. Pulmonary functions after laparoscopic and open cholecystectomy. *Surg Endos Interven Techns.* 2002. 16(1): p. 163-165.

[18] Karayiannakis A. Postoperative pulmonary function after laparoscopic and open cholecystectomy. *British J Anaesthesia.* 1996; 77(4): 448-452.

[19] Kravchenko I, Eberle L, Nesterkina M, Kobernik A. Anti-inflammatory and analgesic activity of ointment based on dense ginger extract (*Zingiber officinale*). *J Herbmед Pharmacol.* 2019; 8(2): 126-132.

[20] Karimi A, Mohammadi-Kamalabadi M, Rafieian-Kopaei M, Amjad L, Salimzadeh I. Determination of antioxidant activity, phenolic contents and antiviral potential of methanol extract of *Euphorbia spinidens* Bornm (Euphorbiaceae). *Trop J Pharmac Res.* 2016; 15(4):759-64.

[21] Vaezi G, Tavasoli Z, S. R-B. Study on the different dosages of *Elaeagnus angustifolia* aqueous extract with and without morphine on the antinociceptive rate in mice. *Research in Medicine. J Shahid Beheshti Uni.* 2011; 35 (1): 27-33.

[22] Froushani SMA, Zarei L, Ghaleh HEG, Motlagh BM. Estragole and methyl-eugenol-free extract of *Artemisia dracunculus* possesses

immunomodulatory effects. *Avicenna J Phytomed.* 2016; 6(5): 526-534.

[23] Abbasi N, Azizi Jalilian F, Abdi M, Saifmanesh M. A comparative study of the antimicrobial effect of *Scrophularia striata* Boiss. Extract and selective antibiotics against *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *J Med Plants* 2007; 6(Suppl 1): 10-18+69.

[24] Bahmani, M., Khaksarian, M., Rafieian-Kopaei, M., Abbasi, N. Overview of the therapeutic effects of *origanum vulgare* and *hypericum perforatum* based on Iran's ethnopharmacological documents. *J Clin Diagn Res.* 2018; 12(7): 1-4.

[25] Shokri Z, Khoshbin M, Koohpayeh A, Abbasi N, Bahmani F, Rafieian-Kopaei, M, Beyranvand F. Thyroid diseases: Pathophysiology and new hopes in treatment with medicinal plants and natural antioxidants. *Int J Green Pharmac.* 2018; 12(3): 473-482.

[26] Abbasi N, Mohammadpour S, Karimi E, Aidi A, Karimi P, Azizi M, et al. Protective effects of *smyrnum cordifolium* boiss essential oil on pentylenetetrazol-induced seizures in mice: Involvement of benzodiazepine and opioid antagonists. *J Biol Regul Homeost Agents.* 2017; 31(3):683-689.

[27] Tajbakhsh M, Karimi A, Tohidpour A, Abbasi N, Fallah F, Akhavan MM. The antimicrobial potential of a new derivative of cathelicidin from *Bungarus fasciatus* against

methicillin-resistant *Staphylococcus aureus*. *J Microbiol*. 2018; 56(2): 128-137.

[28] Razmjoei D, zarei Z, Akbari M. Study of Ethnobotani Some Medicinal Plants in Abadeh, Fars Province. *J Ecophysiol Crops*. 2015; 7 (3): 222-234.

[29] Sharififaa F, MoharamKhani MR, Moaatar F, Babakhanlou P, Khodami M. Ethnobotanical Study of Medicinal Plants of Joopar Mountains of Kerman Province, Iran. *J Kerman Uni Med Sci*. 2014; 21(1): 37-51

[30] Ganjali A, Khaksafidi A. Identification of Some pharmaceutical plant species in Zabol (Iran, Sistan and Baluchistan province) and Their application in traditional medicine . *J Iran Islam Trad Med*. 2015; 6 (1): 89-96.

[31] Khodayari H, Amani SH, Amiri H. Ethnobotanical study of Northeast of Khuzistan province. *Med Plants Ecophytochem J*. 2013; 2(4):12-26.

[32] Mirdalimi SZ, Heshmati GHA, Barani H. Ethnobotanical and ethnoecological survey on medicinal species (case study Kechik Rangelands in the Northeast Golestan Province). *Indig Know*. 2015; 1(2): 129–154.

[33] Khodayari H, Amani SH, Amiri H. Ethnobotanical study of Kichag city. *Herb Med J*. 2013; 7(3):12-26.

[34] Ghavam M, Jeihouni-Naeini H, Kianisalmi S. Indigenous and Traditional Knowledge of the Use of Medicinal Herbs in the Nain County. *Native Iran Knowledge*. 2015; 2(4): 179-204.

[35] Rojni H, Akbari F. Indigenous knowledge of the use of medicinal plants in the village of Erim, the area of the Thousand Jerim of Neka. The first national conference of medicinal herbs and aromatic herbs of Iran, Gonbad Kavoods, May 1, 2017, 2-4.

[36] Ghasemi Pirbalouti A, Momeni M, Bahmani M. ETHNOBOTANICAL STUDY OF MEDICINAL PLANTS USED BY KURD TRIBE IN DEHLORAN AND ABDANAN DISTRICTS, ILAM PROVINCE, IRAN. *Afr J Tradit Complement Altern Med*. 2013 10(2):368-385.

[37] Razmjoue D, Zarei Z, Armand R. Ethnobotanical Study (Identification, Medical Properties and How to Use) of some Medicinal Plants of Behbahan city of Khuzestan Province, Iran. *J Med Plants*. 2018; 16(4): 33-49.

[38] Abedi M., Panahi C., Sattariyan A., Habibi M. Study of Ethnobotane of Medicinal Plants in the Razujorgh County of North Khorasan Province. The first national conference of medicinal herbs and aromatic herbs of Iran, Gonbad Kavoods, May 1, 2017, Available in [conf.gonbad.ac.ir]

[39] Sharififar F, Kouhpayeh A, Motaghi MM, AmirKhosravi A, Pou-Mohseninasab A. The reviews ethnobotany of medicinal plants city of Sirjan, Kerman Province. *J Herbal Drugs*. 2010; 3:19-28.

[40] Ranmanesh M, Najafi SH, Yousefi M. Ethnobotanical study of Medicinal Plants of Sistan region. *J Herbal Drugs*. 2010; 2: 61-8.

- [41] Mozaffari Nejad as, Kamkar A, Giri A, Pourmahmoudi AA. Ethnobotany and folk medicinal uses of major trees and shrubs in Northern Iran. *J Med Plants Res.* 2013; 7(7): 284-289.
- [42] Dolatkahi M, Ghorbani Nahouji M, Mehrafarin A, Amininejad GHR, Dolatkahi A. Ethnobotonic Study of Medicinal Plants in Kazeroon: Identification, Distribution and Traditional Costs. *J Med Plants.* 2012; 11(2): 163-178.
- [43] Safa O, Soltanipoor MA, Rastegar S, Kazemi M, Nourbakhsh Dehkordi KH, Ghannadi A. An ethnobotanical survey on hormozgan province, Iran. *Avicenna J Phytomed.* 2012; 3(1): 64-81
- [44] Procedural sedation in the acute care setting". *Am Fam Physician.* Jan 2005.
- [45] Jamshidi-Kia F, Lorigooini Z, Amini-Khoei H. Medicinal plants: past history and future perspective. *J Herbmed Pharmacol.* 2018; 7(1):1-7.
- [46] Schulz V, Hansel R, Tyler V.E. *Rational Phytotherapy.* SpringerVerlag, Berlin and Heidelberg. 2000; 101-104.
- [47] Bos R, Woerdenbag HJ, Van Putte, PMS, Hendriks H, Scheffer JJC. Seasonal variations of the essential oil, valerenic acid and derivatives, and velepotriates in valeriana officinalis roots and rhizomes, and the selection of plants suitable for phytomedicines. *Planta Med.* 1998; 64: 143–147.
- [48] Vorbach E U, Gortelmeyer R, Bruning J. Therapie von insomnien: wirksameit und vertraglichkeit eines Baldrian-preparates. *Psychopharmakotherapie.* 1996; 3: 109–115
- [49] Ziegler G, Plotch M, Miettinen-Baumann A, Collet W. Efficacy and tolerability of valerian extract LI 156 compared with oxazepam in the treatment of non-organic insomnia – a randomized, double-blind, comparative clinical study. *Eur J Med Res.* 2002; 7: 480–486
- [50] Gholamzadeh S, Zare S, Ilkhanipoor M. Anxiolytic effect of *Echium amoenum* during different treatment courses. *Res J Biolog Sci.* 2008; 3(2):176-8.
- [51] Shafaghi B, Naderi N, Tahmasb L, Kamalinejad M. Anxiolytic effect of *Echium amoenum* L. in mice. *Iranian J Pharmac Res.* 2010; 20:37-41.
- [52] Rabbani M, Sajjadi SE, Vaseghi G, Jafarian A. Anxiolytic effects of *Echium amoenum* on the elevated plus-maze model of anxiety in mice. *Fitoterapia.* 2004; 1; 75(5): 457-64.
- [53] Kyrou I, Christou A, Panagiotakos D, Stefanaki C, Skenderi K, Katsana K, Tsigos C. Effects of a hops (*Humulus lupulus* L.) dry extract supplement on self-reported depression, anxiety and stress levels in apparently healthy young adults: a randomized, placebo-controlled, double-blind, crossover pilot study. *Hormones.* 2017; 16(2):171-80.
- [54] Shishehgar R, Rezaie A, Nazeri M. Study of sedation, pre-anesthetic and anti-anxiety effects of hop (*Humulus lupulus* L.) extract compared with diazepam in rats. *J Animal Vet Advances.* 2012.

[55] Gulec M1, Ozkol H, Selvi Y, Tuluçe Y, Aydin A, Besiroglu L, Ozdemir PG. Oxidative stress in patients with primary insomnia. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012; 1: 37(2): 247.

[56] Bahmani M, Sarrafchi A, Shirzad H, Rafieian-Kopaei M. Autism: Pathophysiology and Promising Herbal Remedies. *Current Pharmaceutical Design*. 2016; 22(3): 277-85.

[57] Shayganni E, Bahmani M, Asgary S, Rafieian-Kopaei M. Inflammaging and cardiovascular disease: Management by medicinal plants. *Phytomed*. 2016; 23(11): 1119-26.

[58] Hosseini Z, Lorigooini Z, Rafieian-Kopaei M, Shirmardi HA, Solati K. A review of botany and pharmacological effect and chemical composition of *Echinophora* species growing in Iran. *Pharmacog Res*. 2017; 9(4): 305–312.

[59] Moradi F, Sewell RDE, Lorigooini Z, Rafieian-Kopaei M. Immunosuppression-lipid Metabolism Interplay and Medicinal Plants in Atherosclerosis: A Review. *Curr Pharmac Design*. 2018; 24:1-5.

[60] Sarrafchi A, Bahmani M, Shirzad H, Rafieian-Kopaei M. Oxidative Stress and Parkinson's Disease: New Hopes in Treatment with Herbal Antioxidants. *Curr Pharmac Design*. 2016; 22(2): 238-46.

[61] Nazarian-Samani Z, Sewell RDE, Lorigooini Z, Rafieian-Kopaei M. Medicinal plants with multiple effects on diabetes mellitus and its

complications: a systematic review. *Curr Diab Rep*. 2018; 18: 72.