


Plant Biotechnology Persa



Online ISSN: 2676-7414

Homepage: https://pbp.medilam.ac.ir

A Comprehensive Review of the Most Important Medicinal Plants Effective in Gestational Diabetes: An In-depth Exploration of Their Pharmacological **Mechanisms**





¹Department of Obstetrics and Gynecology, School of Medicine, Kowsar Woman's General University Hospital, Urmia University of Medical Sciences, Urmia, Iran

Article Info

Article type:

Review Article

Article History:

Received: 16 Dec 2024 Revised: 14 May 2025 Accepted: 29 May 2025 Published Online: 20 Sep 2025

[™] Correspondence to:

Shabnam Vazifekhah

Email:

shabnam.vazifekhah@yahoo.com

ABSTRACT

Objective: Gestational diabetes arises due to the body's inability to effectively regulate blood glucose levels, caused by hormonal changes and insulin resistance, and if left untreated, it jeopardizes the health of both the mother and the fetus. This condition is classified into two types: A1 (manageable through diet) and A2 (requiring medication or insulin). Genetic factors and excess weight can also contribute to its onset. The aim of this study is to provide a comprehensive review of the most significant medicinal plants native to Iran that are effective in managing gestational diabetes and to explore their pharmacological mechanisms.

Methodology: For this review, a search was conducted in reliable databases such as PubMed, Scopus, SID, and Google Scholar, as well as traditional Iranian medicine reference books, to gather articles related to medicinal plants effective against gestational diabetes. The medicinal plants related to gestational diabetes in traditional Iranian medicine were selected for this review. The objective of this study was to identify plants that can be used for managing gestational diabetes with minimal side effects.

Results: In traditional Iranian medicine, several medicinal plants have been recommended for the management of gestational diabetes, including Trachyspermum ammi (L.) Sprague, Foeniculum vulgare Mill., Origanum vulgare L., Myrtus communis L., Eucalyptus globulus Labill., Zingiber officinale Roscoe, Salvia officinalis L., Cinnamomum verum J. Presl, Aloysia citrodora Paláu, Lippia citriodora Kunth, Cucurbita pepo L., Artemisia absinthium L., Artemisia herba-alba Asso, Juglans regia L., Cuminum cyminum L., Vicia faba L., Olea europaea L., Thymus vulgaris L., Mentha piperita L., Cuminum cyminum L., Rosmarinus officinalis L., Solanum melongena L.. These plants have been used for controlling gestational diabetes in traditional Iranian medicine.

Conclusion: Gestational diabetes, as a common condition during pregnancy, requires careful and effective management to prevent serious complications for both the mother and the fetus. In traditional Iranian medicine, many medicinal plants are recognized as adjunctive treatments for managing gestational diabetes. The use of these plants could serve as a natural and costeffective strategy alongside standard medical treatments, particularly in cases of gestational diabetes. However, it is crucial that the consumption of these plants be supervised by a healthcare professional to avoid any potential side effects or drug interactions.

Gestational Diabetes, Medicinal Plants, Pharmacological Mechanisms, Blood Sugar Control, Insulin Resistance

How to cite this paper

Vazifekhah SH. A Comprehensive Review of the Most Important Medicinal Plants Effective in Gestational Diabetes: An In-depth Exploration of Their Pharmacological Mechanisms. Plant Biotechnology Persa. 2025; 7(4): 57-64. DOI: 10.61882/pbp.7.4.8



Introduction

Insulin resistance and hyperglycemia are more common in pregnant women [1]. Insulin is a hormone that converts blood sugar into glycogen and stores it in cells [2]. During pregnancy, hormones secreted by the placenta and fetus reduce the body's sensitivity to insulin. In most cases, the body compensates for this by increasing insulin production. However, in some women, insulin production is insufficient, and blood sugar levels rise. This condition is known as gestational diabetes [3]. In some pregnant women, blood sugar levels increase in the middle of pregnancy, leading to gestational diabetes. This complication typically occurs between the 24th and 28th weeks of pregnancy. According to CDC reports, the prevalence of gestational diabetes is estimated to be between 2% and 14% [14]. During pregnancy, hormonal changes can reduce insulin effectiveness and increase blood sugar, which can result in gestational diabetes in some women. Risk factors include excess weight, family history of diabetes, and the use of certain medications [5]. Diabetes, as a condition of multifactorial origin including genetic and environmental factors, accounts for 3.5% of the mortality cases because of non-communicable chronic [6]. Gestational diabetes is usually asymptomatic, but in some cases, mild symptoms such as excessive thirst, frequent urination, dry mouth, fatigue, blurred vision, and genital itching or infections may occur [7]. Risk factors for gestational diabetes include family history, overweight, and high-risk pregnancies. Early diagnosis is essential to prevent maternal and fetal complications [8]. diabetes Untreated gestational can have significant consequences for the baby, including macrosomia (high birth weight), increased risk of respiratory disorders such as respiratory distress syndrome, neonatal hypoglycemia, and a higher likelihood of obesity and type 2 diabetes later in life [9]. Gestational diabetes is typically diagnosed between the 24th and 28th weeks of pregnancy, and women with risk factors should be screened before the 15th week. Diagnosis is made through a two-step blood glucose test after glucose consumption [10].

A suitable diet for gestational diabetes includes a balanced intake of proteins, complex carbohydrates, healthy fats, and fiber [11]. This diet should be consumed regularly in small meals throughout the day to properly control blood sugar [12]. Processed foods and sugary beverages should be avoided to reduce the risks associated with gestational diabetes [13]. While gestational diabetes cannot be fully prevented, the risk of developing it can be reduced by maintaining healthy habits such as consuming fiber-rich, lowfat foods and engaging in regular physical activity. Maintaining a healthy weight before pregnancy and consulting with a healthcare provider about appropriate weight gain during pregnancy can also prevent gestational diabetes and type 2 diabetes [14]. Gestational diabetes is a condition characterized by elevated blood sugar during pregnancy, and its treatment includes a healthy diet, exercise, and in some cases, insulin therapy. Blood sugar control and monitoring fetal health are essential to prevent serious complications such as preterm birth and high birth weight [15]. Traditional Iranian Medicine, with its deep-rooted history, has long advocated the use of medicinal herbs for treating various ailments [16]. In medicine, medicinal plants traditional recommended for managing gestational diabetes to help lower blood sugar levels [17]. Dietary modifications, including the consumption of vegetables, whole grains, and natural antiinflammatory foods, combined with mild exercise, can aid in blood sugar control and improve the pregnant woman's condition [18]. Green tea, cinnamon, bitter melon, and other medicinal plants have been shown to be effective in reducing blood sugar and controlling gestational diabetes. These plants can be consumed as teas, powders, or supplements to help regulate blood sugar levels [19]. This study aims to provide a comprehensive review of the medicinal plants native to Iran that are effective in managing gestational diabetes and explore their pharmacological mechanisms.

Methodology

In this review, a search was conducted in reputable databases including PubMed, Scopus, SID, and Google Scholar, as well as in reference books of traditional Iranian medicine, to gather articles related to medicinal plants effective in treating gestational diabetes. Medicinal plants known in traditional Iranian medicine for controlling, managing, and treating gestational diabetes were selected for an in-depth review. The inclusion criteria for this study included articles and resources that specifically address the medicinal plants native to Iran and their effects on treating gestational diabetes. Conversely, sources with insufficient or unrelated information were excluded from the study.

Results

Traditional Iranian medicine uses various medicinal plants to control gestational diabetes, helping regulate blood sugar levels and improving the health of both the mother and the fetus. Notable plants include *Trachyspermum ammi* (L.) Sprague, Foeniculum vulgare Mill., Origanum vulgare L., Myrtus communis L., Eucalyptus globulus Labill., Zingiber officinale Roscoe, Salvia officinalis L., Cinnamomum verum J. Presl, Aloysia citrodora Paláu, Lippia citriodora Cucurbita pepo L., Artemisia absinthium L., Artemisia herba-alba Asso, Juglans regia L., Cuminum cyminum L., Vicia faba L., Olea europaea L., Thymus vulgaris L., Mentha piperita L., Cuminum cyminum L., Rosmarinus officinalis L., Solanum melongena L.. These plants have been used in traditional Iranian medicine to manage gestational diabetes and are employed as natural treatments to reduce the complications of the condition. Further results on medicinal plants used for managing gestational diabetes in traditional Iranian medicine are detailed in Table 1.

Table 1: Medicinal Plants for Gestational Diabetes [17-31]

| Persian Name | Scientific Name | Plant Family | Used Part | Traditional Use | Mechanism of Action |
|-------------------|---|---------------|---------------|-----------------------------------|--|
| Zanayan | <i>Trachyspermum ammi</i> (L.) Sprague | Apiaceae | Seed | Consumed as tea or extract | Antimicrobial and anti-inflammatory effects |
| Fennel | Foeniculum vulgare Mill. | Fabaceae | Seed | Consumed as tea or extract | Boosts immune system and anti- inflammatory effects |
| Shirazi Thyme | <i>Origanum vulgare</i> L. | Lamiaceae | Leaf | Consumed as tea or extract | Antimicrobial and inflammation reduction |
| Myrtle | <i>Myrtus communis</i> L. | Myrtaceae | Leaf | Consumed as leaf extract or tea | Boosts immune system and antimicrobial properties |
| Eucalyptus | Eucalyptus globulus Labill. | Lauraceae | Leaf | Consumed as decoction or extract | Antimicrobial and inflammation reduction |
| Ginger | Zingiber officinale Roscoe | Zingiberaceae | Root | Consumed as decoction or powder | Antimicrobial and anti-inflammatory effects |
| Sage | <i>Salvia officinalis</i> L. | Lamiaceae | Leaf | Consumed as tea or extract | Antimicrobial properties and boosts immune system |
| Cinnamon | Cinnamomum verum J. Presl | Lauraceae | Bark | Consumed as decoction or powder | Antimicrobial and anti-inflammatory effects |
| Lemon Verbena | <i>Aloysia citrodora</i> Paláu | Verbenaceae | Leaf | Consumed as tea or extract | Antimicrobial and inflammation reduction |
| Lemon Balm | <i>Lippia citriodora</i> Kunth | Verbenaceae | Leaf | Consumed as tea or extract | Boosts immune system and antimicrobial properties |
| Pumpkin | <i>Cucurbita pepo</i> L. | Cucurbitaceae | Fruit | Consumed raw or dried | Reduces inflammation and boosts immune system |
| Wormwood | <i>Artemisia absinthium</i> L. | Asteraceae | Leaf and root | Consumed as decoction or extract | Antimicrobial and anti-inflammatory effects |
| White Wormwood | <i>Artemisia herba-alba</i> Asso | Asteraceae | Leaf | Consumed as decoction or powder | Reduces inflammation and boosts immune system |
| Walnut | <i>Juglans regia</i> L. | Juglandaceae | Fruit | Consumed as tea or powdered seeds | Antimicrobial and anti-inflammatory effects |

Shabnam Vazifekhah

| Black Cumin | Cuminum cyminum L. | Apiaceae | Seed | Consumed as tea or | Antimicrobial properties and boosts |
|-------------|-----------------------------|------------|-------|----------------------------|-------------------------------------|
| | | | | powdered seeds | immune system |
| Broad Bean | <i>Vicia faba</i> L. | Fabaceae | Seed | Consumed as tea or | Boosts immune system and |
| | | | | powdered seeds | antimicrobial properties |
| Olive | <i>Olea europaea</i> L. | Oleaceae | Fruit | Consumed raw or dried | Antimicrobial and anti-inflammatory |
| | | | | | effects |
| Thyme | <i>Thymus vulgaris</i> L. | Lamiaceae | Leaf | Consumed as tea or extract | Boosts immune system and |
| | | | | | antimicrobial properties |
| Mint | <i>Mentha piperita</i> L. | Lamiaceae | Leaf | Consumed as decoction or | Antimicrobial and anti-inflammatory |
| | | | | extract | effects |
| Zanayan | Cuminum cyminum L. | Apiaceae | Seed | Consumed as tea or extract | Antimicrobial properties and boosts |
| (Again) | | | | | immune system |
| Rosemary | Rosmarinus officinalis L. | Lamiaceae | Leaf | Consumed as decoction or | Antimicrobial and inflammation |
| | | | | extract | reduction |
| Eggplant | <i>Solanum melongena</i> L. | Solanaceae | Fruit | Consumed raw or dried | Antimicrobial and anti-inflammatory |
| | | | | | effects |

Discussion

Diabetes is a metabolic disorder characterized by impaired insulin production or function, leading to elevated blood glucose levels. Gestational diabetes is a form of diabetes that develops during pregnancy and typically resolves after childbirth. However, it can pose risks to both the mother and the baby. Proper control and timely medical care can help prevent serious complications associated with this condition [35].

Research indicates that over 1200 plant species with antidiabetic properties have been identified. The most significant active compounds found in these plants include imidazolines, alpha-lipoic acid, flavonoids, and saponins. The mechanisms of action of these compounds involve stimulating insulin secretion, antioxidant activity, enzyme inhibition, and the suppression of glucose biosynthesis [36]. A review study exploring the use of medicinal plants in diabetes management highlights several plants, such as Allium cepa, Allium sativum, Gymnema sylvestre, and Silybum marianum, as effective treatments. The study also emphasizes the potential risks of using these plants unwittingly, such as drug interactions and fluctuations in blood glucose levels [37]. Another investigation examines the role of plants such as garlic, onion, fenugreek, milk thistle, and green tea in managing diabetes. It underscores the importance of cautious usage to prevent adverse drug interactions and challenges in blood sugar control. The study also points to the significance of dietary changes and alternative treatments in preventing and managing Type 2 diabetes and assisting in the treatment of Type 1 diabetes [38].

A review of clinical trials on medicinal plants for diabetes treatment in Iran from 2010 to 2020 reveals that certain plants, including cinnamon and cumin, have shown positive effects on blood glucose levels and HbA1c [39]. Several medicinal plants, such as *Apium graveolens*, *Alyssum desertorum*, *Arctium lappa*, *Avena sativa*, *Berberis integerima*, *Cerasus microcarpa*, *Crataegus aronia*, *Allium schoenoprasum*, *Urtica dioica*, *Phlomis aucheri*, *Salvia*

aethiopis, Melilotus officinalis, Tragopogon pratensis, Euphorbia helioscopia, Salvia officinalis, and Salix aegyptiaca, have been used in West Azerbaijan, East Azerbaijan, and Ardabil provinces for treating diabetes. These plants have long been employed in traditional medicine in these regions due to their anti-diabetic properties [40].

Further research has reported that 282 plant species from 63 families have been documented for treating diabetes in Iran. Urtica dioica, along with Citrullus colocynthis, Teucrium polium, Juglans regia, and Trigonella foenumgraecum, are among the most frequently used plants in diabetes treatment. Phylogenetic relationships between the plant families and their parts used for diabetes treatment have shown significant correlations based on evolutionary plant patterns [41]. A total of 85 studies (18 human trials and 67 animal studies) have examined 62 plants, with randomized controlled trials (RCTs) assessing the quality of studies using the Jadad scale. Among the RCTs, the best results in blood sugar control were found with Aloe vera, Citrullus colocynthis, Plantago ovata, Silybum marianum, Rheum ribes, and Urtica dioica [42]. In Urmia, medicinal plants such as Achillea millefolium L., Alyssum desertorum Stapf., Arctium lappa L., Avena sativa L., Berberis integerima Bunge., Cerasus microcarpa, Cinnamomum verum, Citrullus colocynthis (L.) Schrad., Crataegus aronia (L.) Bosc ex Dc., Juglans regia, L. album L., and Sanguisorba *minor* Scop. are utilized for treating diabetes [43]. Medicinal plants such as Nigella sativa, Aconitum napellus, Agaricus albolutescens, and Euphorbia cedrorum, among others, are used for gestational diabetes [44]. In many diseases and disorders [45-49], turning to nature and embracing traditional or natural therapeutic approaches can serve as a beneficial and complementary strategy helping to alleviate symptoms, support overall well-being, and enhance patients' quality of life [50].

Conclusion

Gestational diabetes is a common concern during pregnancy that requires careful monitoring and treatment to prevent serious complications for both the mother and the fetus. In traditional Iranian medicine, numerous medicinal plants are recognized as complementary treatments for managing gestational diabetes. These plants can be beneficial as natural and cost-effective alternatives alongside conventional medical treatments, especially in the case of gestational diabetes. However, the use of these plants should be supervised by a qualified healthcare provider to prevent potential side effects and interactions with other medications.

Statements and Declarations Funding support

The authors did not receive support from any organization for the submitted work.

Competing interests

The authors have no competing interests to declare that are relevant to the content of this article.

Ethics approval

This study was performed in line with the principles of the Declaration of Helsinki.

Acknowledgments

The authors would like to express their gratitude to the clinical research development unit of Imam Khomeini Hospital, Urmia University of Medical Sciences, for English editing.

References

- Boroumand F, Ghayur S, Gharaaghaji R, Vazifekhah S. Efficacy of prophylactic use of metformin in prevention of gestational diabetes mellitus in nondiabetic obese pregnant women. J Obstet Gynecol Cancer Res. 2022;7(6):524-9.
- 2. McIntyre HD, Catalano P, Zhang C, Desoye G, Mathiesen ER, Damm P. Gestational diabetes mellitus. Nat Rev Dis Primers. 2019;5(1):47. doi: 10.1038/s41572-019-0098-8.
- Buchanan TA, Xiang AH. Gestational diabetes mellitus. J Clin Invest. 2005;115(3):485–91.

- 4. Coustan DR. Gestational diabetes mellitus. Clin Chem. 2013;59(9):1310-21.
- 5. Buchanan TA, Xiang A, Kjos SL, Watanabe R. What is gestational diabetes?. Diabetes Care. 2007;30(Suppl 2): 105. doi: 10.2337/dc07-s201.
- Rad AH, Abbasalizadeh S, Vazifekhah S, Abbasalizadeh F, Hassanalilou T, Bastani P, et al. The future of diabetes management by healthy probiotic microorganisms. Curr Diabetes Rev. 2017;13(6):582-9.
- 7. Ryan EA. Diagnosing gestational diabetes. Diabetologia. 2011;54(3):480–6. doi: 10.1007/s00125-010-2005-4.
- Kampmann U, Madsen LR, Skajaa GO, Iversen DS, Moeller N, Ovesen P. Gestational diabetes: a clinical update. World J Diabetes. 2015;6(8):1065. doi: 10.4239/wjd.v6.i8.1065.
- 9. Kjos SL, Buchanan TA. Gestational diabetes mellitus. N Engl J Med. 1999;341(23):1749–56.
- 10. Kjos SL, Buchanan TA. Gestational diabetes mellitus. N Engl J Med. 1999;341(23):1749–56.
- 11. Reece EA, Leguizamón G, Wiznitzer A. Gestational diabetes: the need for a common ground. Lancet. 2009;373(9677):1789–97.
- 12. Mecacci F, Lisi F, Vannuccini S, Ottanelli S, Rambaldi MP, Serena C, et al. Different gestational diabetes phenotypes: which insulin regimen fits better?. Front Endocrinol. 2021;12:630903. https://doi.org/10.3389/fendo.2021.630903
- 13. Persson B, Stangenberg M, Hansson U, Nordlander E. Gestational diabetes mellitus (GDM): comparative evaluation of two treatment regimens, diet versus insulin and diet. Diabetes. 1985;34(Suppl 2):101–5.
- 14. Cheung NW. The management of gestational diabetes. Vasc Health Risk Manag. 2009;5:153-64.
- 15. Xu YX, Xi S, Qian X. Evaluating traditional Chinese medicine and herbal products for the treatment of gestational diabetes mellitus. J Diabetes Res. 2019;2019:9182595.
- 16. Vazifekhah S. The Most Effective Medicinal Plants for Toxoplasmosis-Induced Miscarriage and of Their Mechanisms of Action. Plant Biotechnol Persa. 2025;7(3):0-0.
- 17. Yessoufou A, Gbenou J, Grissa O, Hichami A, Simonin AM, Tabka Z, et al. Anti-hyperglycemic effects of three medicinal plants in diabetic pregnancy: modulation of T cell proliferation. BMC Complement Altern Med. 2013;13:1–3. doi: 10.1186/1472-6882-13-77.
- 18. Arista DM, Amelia R, Fitriani D, Khotimah H, Ratnaningrum SD, Irwanto Y, et al. Gestational Diabetes Mellitus: An overview and its potential treatment with herbs. GSC Biol Pharm Sci. 2023;23(3):261–73.
- 19. Bernstein N, Akram M, Yaniv-Bachrach Z, Daniyal M. Is it safe to consume traditional medicinal plants during pregnancy?. Phytother Res. 2021;35(4):1908–24. doi: 10.1002/ptr.6935.
- 20. Aghili Khorasani MH. Qarabadin-e-Kabir. Tehran: Research Institute for Islamic and Complementary Medicine, Tehran University of Medical Sciences; 2009.

- 21. Avicenna. The Canon of Medicine (Al-Qanun fi al-Tibb). Translated by Laleh Bakhtiar. Chicago: Kazi Publications; 1999.
- 22. Arzani MA. Tebb-e-Akbari. Tehran: Research Institute for Islamic and Complementary Medicine, Tehran University of Medical Sciences; 2008.
- 23. Momen Tonekaboni H. Tohfat al-Momenin. Tehran: Nashr-e Shahr; 2007.
- 24. Zargari A. Medicinal Plants. Tehran: Tehran University Press; 1992.
- 25. Jorjani SE. Zakhireh Kharazmshahi (Treasure of Kharazm Shah). Tehran: Iranian Academy of Medical Sciences; 2001.
- 26. Ghaznavi H. Traditional Iranian Medicine and Medicinal Plants. Tehran: Nashr-e Elm; 2011.
- 27. Khorasani M, Najmabadi M. Medicinal Plants of Iran. Tehran: Tehran University Press; 2006.
- 28. Sadr E. Encyclopedia of Medicinal Plants. Tehran: Jangal Publishing House; 2010.
- 29. Mir Heidar H. Plant Sciences and Medicinal Herbs of Iran. Tehran: Islamic Culture Press; 1993.
- 30. Nazem Javedan M. Kholasat al-Hikmah (Summary of Wisdom). Tehran: Institute of Traditional Medicine and Materia Medica Research; 2012.
- 31. Dehkhoda A. Dehkhoda Dictionary: Plant Names. Tehran: Tehran University Press; 1998.
- 32. Davari Ardekani A. Principles of Iranian Traditional Medicine. Tehran: University of Tehran Press; 2015.
- 33. Ahvazi A. Kamel al-Sanaat al-Tibbiyya. Tehran: Academy of Medical Sciences of the Islamic Republic of Iran: 2008.
- 34. Shams Ardakani M, Ghorbani A. Herbal Medicines in Iranian Traditional Medicine. Tehran: Medical Sciences University Press; 2015.
- 35. Incomplete entry: Please provide full citation for "32".
- 36. Bat'haei SZ, Makarezadeh N, Shirali S. A review on the mechanisms of action of plant bioactive compounds in the treatment of diabetes mellitus. Herbal Plants. 2012;11(44):1–24. Available from: https://sid.ir/paper/15171/fajv
- 37. Fallah Huseini H, Fakhrzadeh H, Larijani B, Shikh Samani A. Review of anti-diabetic medicinal plant used in traditional medicine. J Med Plants. 2006;5(S2):1–8. Available from: http://jmp.ir/article-1-583-en.html
- 38. Fallah Hosseini H, Fakhrzadeh H, Ardestani Larijani MB, Sheikh Samani AH. A review of medicinal plants used in diabetes. Herbal Plants. 2005;5(Special Issue on Diabetes):1–8. Available from: https://sid.ir/paper/15205/fa
- 39. Fallah B, Mehrabbeik A, Salmani N. A Review of Clinical Trial Studies on Medicinal Plants in the Treatment of Diabetes. Int J Diabetes Obes. 2022;14(2):117–30.
- 40. Mohammadi E, Abdi F. Medicinal Plants Effective on Diabetes in Northwest of Iran. Plant Biotechnol Persa. 2025;7(1):4–9.
- 41. Labbafi M, Mohammadi T, Babaei A, Abbasi S, Avazzadeh A, Khalaj H, et al. An overview of documented medicinal plants used for the treatment of diabetes in Iran with ethnobotanical and evolutionary perspective (2002–2022). Ethnobot Res Appl.

- 2024;29:1–53. Available from: https://ethnobotanyjournal.org/index.php/era/article/view/6387
- 42. Rashidi AA, Mirhashemi SM, Taghizadeh M, Sarkhail P. Iranian medicinal plants for diabetes mellitus: a systematic review. Pak J Biol Sci. 2013;16(9):401–11. doi: 10.3923/pjbs.2013.401.411
- 43. Bahmani M, Zargaran A, Rafieian-Kopaei M, Saki K. [Title of the article]. Asian Pac J Trop Med. 2014;7(Suppl 1): 348–S354.
- 44. Yedjou CG. The management of diabetes mellitus using medicinal plants and vitamins. Int J Mol Sci. 2023;24(10):9085. doi: 10.3390/ijms24109085.
- 45. Mahmud Hussen B, Noori M, Sayad B, Ebadi Fard Azar M, Sadri Nahand J, Bayat M, Babaei F, Karampour R, Bokharaei-Salim F, Mirzaei H, Moghoofei M. New potential MicroRNA biomarkers in human immunodeficiency virus elite controllers, human immunodeficiency virus infections, and coinfections with hepatitis B virus or hepatitis C virus. Intervirology. 2023 Dec 20;66(1):122–135.
- 46. Ghanbari A, Nouri M, Darvishi M. Evaluation of relationship between serum hemoglobin A1C level and severity of diabetic foot ulcers based on Wagner criteria. J Med Chem Sci. 2023;6:2234–2241.
- 47. Darvishi M, Nouri M, Zahir M, Asli M, Hejripoor SZ, Karimi E. Overview of human papillomavirus infection. Infect Disord Drug Targets. 2024 Mar 1;24(2):65–76.
- 48. Nouri M, Kamakifar P, Khodabandehlou N, Nahand JS, Tavakoli A, Norooznezhad F, Sorayyayi S, Babaei F, Mostafaei S, Moghoofei M. Association between Parvovirus B19 and anemia in HIV-infected patients. Med J Islam Repub Iran. 2019 Dec 16;33:137.
- 49. Darvishi M, Noori M, Nazer MR, Soleiman-Meigooni S, Forootan M. The relationship between Helicobacter pylori and extra-gastrointestinal infections. Iran J Med Microbiol. 2020 Nov 10;14(6):543–565.
- 50. Darvishi M, Hashemi Rafsanjani SMR, Nouri M, Abbaszadeh S, Heidari-Soureshjani S, Kasiri K, Rahimian G. Biological mechanisms of polyphenols against Clostridium difficile: A systematic review. Infect Disord Drug Targets. 2025 May;25(3): 18715265313944. doi: 10.2174/0118715265313944240726115600