

Plant Biotechnology Persa



Online ISSN: 2676-7414

Homepage: https://pbp.medilam.ac.ir

Effects of Using Salicylic Acid, Potassium Chloride, and a Mixture of Salicylic Acid and Potassium Chloride Treatments on Salt-stressed Cucumis Sativus CV.

Malaysia Timun2 Germination

Samar Jasim Mohammed 1 , Rosimah Nulit 2 , and Mohamed I. A. Fayed 3 D

¹Biology Department, College of Science, University of Misan, Amarah, Iraq; Email: Samarjasim@uomisan.edu.iq
²Biology Department, Faculty of Science, University of Putra Malaysia, Malaysia; Email: rosimahn@upm.edu.my
³Ph.D. Agricultural Engineering, Faculty of Agriculture, Zagazig University, Egypt; Email: dr_eng.fayed@yahoo.com
Corresponding Author, Biology Department, College of Science, University of Misan, Amarah, Iraq; Email: Samarjasim@uomisan.edu.iq

Article Info Abstract

Article type: Objective: In constructing a liquid enhancer, 300 mM NaCl was primed for 72 hours on sterile

Original Article MTi2 seeds.

Methods: After that, deionized water was used as a control, and SA alone (salicylic acid) (0.25, 0.5, 0.75, 1 mM) and only KCl (Potassium chloride) (10, 20, 30, 40, and 50 mM) were applied. As previously stated, germination parameters were computed. Then, the appropriate ratio of KCl to SA was combined, and its efficacy as a germination activator on the Salt-stressed MTi2 seeds was

examined. The data analysis software used was SPSS Windows version 22.

Results: To find the significant difference between treatments, data are first subjected to a two-way ANOVA with p \le 0.05 confidence level. For purpose of comparing means, DMRT is next applied at a p \le 0.05. According to the results, the best concentrations for boosting the germination and early growth of MTi2 seedlings in comparison to the control treatment were found to be 20–30 mM KCl and 0.5–0.75 mM SA. Furthermore, MTi2 seedling germination and early growth were more than 1x higher when the best concentrations of KCl (20-30 mM) and SA (0.5-0.75 mM) were

combined.

Conclusion: Salicylic acid (SA) and low levels of KCl applied to salt-stressed MTi2 seeds can help reduce the negative effects of salinity stress and enhance the percentage, rate, vigour, length, and biomass of the seedlings that germinate. As the conclusion, salt-stressed MTi2 seeds can benefit from an enhancer that increases germination at low concentrations of KCl and SA.

Keywords:

December 2023

Article History:

20 November 2023

2023

2023

Received: 02 September

Received in revised form:

Accepted: 23 December

Published online: 31

Germination, Salinity stress, Seedlings, Salicylic acid, Potassium chloride

Plant Biotechnology Persa 2022; 5(2): 56-67.



DOI: 10.61186/pbp.5.2.7

Introduction

Salinity affects the pH and availability of nutrients in soils, such as manganese, iron, and phosphorus [11]. Though it came from South Asia, cucumbers may now be found on most continents. Saline groundwater in semi-arid regions makes this plant a valuable crop for greenhouses. Consequently, further research is required to determine how salt affects this plant's ability to germinate [12].

It was determined that the cucumber plant was a glycophyte, meaning that it was somewhat susceptible to salinity. Previous research has demonstrated that salicylic acid (SA) plays a significant role in regulating how plants respond to a variety of abiotic stressors, including salt and water stress [13, 14]. Shakirova [15]; and Srivastava [16] discovered that it is crucial to the plant species' defense system against stress. Many application techniques were employed to protection different plant species in response to abioticstresses like salt, including soaking seeds in salicylic acid, Spraying or irrigating with SA solution after adding SA to the hydroponic solution [17, 18].

This study was conducted on the Malaysia cucumber cv. MTi2 which is the best known and most popular cucumber cultivars among the locals. Cucumber (Cucumis sativus) belongs to the gourd, family Cucurbitaceae. It is a widely cultivated, creeping vine that bears cylindrical, fruits that are used as culinary vegetables [19]. The cucumber originated from South Asia, but is currently found on most continents. This plant is an important greenhouse crop in semi- arid areas with saline ground water.

The present study aims to develop a liquid enhancer that will help salt-stressed *Cucumber sativus* cv. MTi2 seeds germinate more efficiently.

Materials and Methodology Seed Materials and Sterilization

Cucumber (Cucumber sativus cv. MTi2) seeds were purchased from MARDI (Malaysia Agriculture Research, and Development Institute, Serdang, Selangor). Sterilization of seeds was done using the technique revealed by Panuccio [20] with a few slight modifications. The cucumber seeds that were chosen were mature, healthy, and of uniform size. They were then surface sterilized for 20 minutes using 5% sodium hypochlorite. The seeds were subsequently cleaned three times using distilled water, that had been sterilized to prevent fungal infections.

Experimental Site

The research project was carried out from June 2015 to October 2016 at the Tissue Culture Laboratory, Department of Biology, University of Putra Malaysia.

Designing an experiment to improve the germination of MTi2 seeds salt-stressed

In this research, salt-stressed MTi2 seedlings were treated with SA (salicylic acid), KCl (potassium chloride), or a combination of both. To do this, initially, the experiment was carried out with SA to determine the optimum concentration of SA that promotes the germination of MTi2 seedlings under salt stress. Second, to determine the optimum KCl concentration that promotes the germination of salt-stressed MTi2 seeds, MTi2 seeds that have been exposed to salt will be germinated in a range of KCl concentrations. Finally, the optimum ratio of SA to KCl will be combined to

determine the ratio that optimizes the germination of MTi2 seeds salt-stressed [21].

Salted Solutions Preparation

Five different concentrations of KCl, which are 10, 20, 30, 40, and 50mM were prepared and deionized water as a control. Four salicylic acid concentrations of 0.25, 0.50, 0.75, and 1 mM were prepared and deionized water was used as a control [21].

Halopriming of MTi2 seeds in NaCl

After being sterilized with MTi2, the seeds were soaked in 50 ml of 300 mM NaCl, and incubated for 72 hrs at 25 °C, in a growth incubator, according to the technique by Afzal [22]; and Elouaer [23].

MTi2 seeds Salt-stressed treated with KCl, SA, and a combination of SA and KCl

Where: GP% is the Germination percentage, NGS is the Number of germinated seeds, TNSS is the Total number of seed sown., GR is the Germination rate, D1 is the Day of the first count, DL is the Day of the last count., SV is the Seed Vigor, LH is the Length of hypocotyl, and LR is the length of radical.

Improvement of salt-stressed MTi2 seed germination using KCl

Seven sterilized MTi2 seeds were placed in two layers of Whatman filter paper No. 1 in separate sterilized Petri dishes (9 cm diameter) containing 5 ml of different concentrations of KCl (10, 20, 30, 40, 50 mM) and as a control, deionized water. All treatment was kept in the growth chamber at 25 ± 1 °C and will be monitored daily until 8 days. Measurements were made on seed vigor,

Improvement of salt-stressed MTi2 seed germination using SA:

Seven sterilized MTi2 seeds were placed on two layers of Whatman filter paper No. 1 in separate sterilized Petri dishes (9 cm diameter) containing 5 ml of different concentrations SA (0.25, 0.5, 0.75, 1 mM) and deionized water as a control. All treatments were kept in the growth chamber at $25 \pm 1^{\circ}$ C and will be monitored daily until 8 days. Measurements were made on seed vigor, germination rate, germination percentage, and early seedling growth [21].

Germination parameters were calculated as follows:

$$GP\% = \frac{NGS}{TNSS} \times 100 [24, 25]$$

$$GR = \frac{NGS}{D1} + \dots + \dots + \frac{NGS}{DL} [26]$$

$$SV = \frac{(LH + LR) \times GP\%}{100} [27]$$

germination rate, germination percentage, and early seedling growth as described in subtitle 2.3.i

Three seedlings from nine replicates of each treatment were selected randomly. The length of seedlings, radicals, and hypocotyls were measured on day 8. For biomass measurement, samples were placed in the oven at 60°C for several days until constant weight was obtained.

Improvement of salt-stressed MTi2 seed germination using a SA and KCl mixture

Seven sterilized MTi2 seeds were placed on two layers of Whatman filter paper No. 1 in separate sterilized Petri dishes (9 cm diameter) comprising the optimal ratio of KCl to SA, as shown in Table 1.

Table 1. Salt-stressed MTi2 seeds germinate using a combination of SA and KCl

Treatment	Concentration (mM)	
Control	Deionized water	
SA (Salicylic acid) + KCl (Potassium chloride)	0.50 SA + 20 KCl	
SA (Salicylic acid) + KCl (Potassium chloride)	0.75 SA + 20KCl	
SA (Salicylic acid) + KCl (Potassium chloride)	0.50 SA + 30KCl	
SA (Salicylic acid) + KCl (Potassium chloride)	0.75 SA + 30KCl	

All treatment was stored in the growing chamber at 25 ± 1 °C and will be monitored daily until 8 days. Measurements were made on seed vigor, germination rate, germination percentage, and early seedling growth as described in subtitle 2.3.i.

Data Analysis

salicylic acid concentrations

SPSS Windows Version 22 was used to conduct the statistical analysis. To investigate the differences in germination response across salt treatments, At a confidence level of p=0.05, a two-way analysis of variance (ANOVA) was conducted. Tukey HSD was then used at p=0.05 for mean comparison. To determine the significance difference utilizing SA, KCl, and a combination of SA and KCl, one-way ANOVA was performed at a confidence level of p=0.05. Tukey HSD was then used at p=0.05 for mean analysis.

Results increased 6X higher in 0.5 mM SA. **Table 2.** Salt-stressed MTi2 seeds germination percentage, germination rate, and seed vigor at different

Conc. (mM)	GP (%)	GR	SV
0	37.1a±7.3	6.0a ±1. 4	2.91a±0.7
0.25	54.3ab±13.1	8.5ab ±2. 2	6.33ab±2.0
0.50	77.1b±7.3	13.2b ±1.4	12.41c±1.7
0.75	80.0b±3.5	12.6b ±0.2	10.87bc±0.7
1	60.0ab±8.3	10.4ab±1.5	5.24a±0.9

Treatment of MTi2 Seeds Salt-Stressed with SA:

The percentage and rate of germination, and Seed Vigor:

Results found that SA (salicylic acid) significantly increased the percentage and rate of germination of salt-stress MTi2 seeds. **Table 2** shows that the percentage of germination is significantly higher in all concentrations of SA than in deionized water.

The highest percentage of germination in 0.75 mM SA. Salicylic acid in all concentrations also increased the speed of germination of salt-stressed MTi2 seeds. However, the highest rate of germination was found in 0.5 mM SA.

The vigor of salt-stressed MTi2 seeds germinated in SA also increased. The vigor of seed is more than 2x higher in SA compared with deionized water. The Vigor of the seed increased 6X higher in 0.5 mM SA.

The mean is used, and five replicates are used to calculate the standard errors of measurement. Significant variations between the means are indicated by superscripts with distinct letters inside the means of each column (a–b) (Tukey HSD test, p<0.05, Appendix 10).

Impact of Salinity on MTi2 Seedlings Grown in Early Stages:

Under salt stress, MTi2 seedlings grow much faster when salicylic acid is applied at all

concentrations (ANOVA, p<0.05) as shown in **Table 3**. The highest length of hypocotyls, radical, seedling length, and biomass was found in 0.50 mM SA, which is 2x higher than in the control.

Table 3. Salicylic acid concentration-dependent measurements of the hypocotyl, radical, seedling, and biomass of salt-stressed MTi2 seeds

Conc. (mM)	HL (cm)	RL (cm)	SL (cm)	Biomass (g)
0	4.1ª±0.4	3.4 ^a ±0.3	7.5°±0.6	0.011 ^a ±0.0021
0.25	$6.0^{ab} \pm 0.6$	$4.7^{ab}\pm0.9$	$10.7^{ab} \pm 1.2$	$0.016^{ab} \pm 0.0024$
0.5	$7.6^{b}\pm0.2$	$8.4^{\circ}\pm0.7$	$16^{c}\pm0.7$	$0.022^{b}\pm0.0016$
0.75	$6.7^{b}\pm0.8$	$6.9^{bc}\pm0.3$	13.6 ^{bc} ±0.6	$0.020^{b}\pm0.0013$
1	$3.8^{a}\pm0.5$	$4.8^{ab}\pm0.6$	$8.6^{a}\pm0.7$	$0.014^{ab}\pm0.0021$

The mean is used, and five replicates are used to calculate the standard errors of measurement. Significant differences between the means are shown by superscripts with distinct letters inside the means of each column (a-c) (Tukey HSD test, p<0.05, Appendix 13-16).

Application of KCl to MTi2 Seeds Under Salt-Stress:

The percentage and rate of germination, and Seed Vigor:

The results found that KCl significantly increased the germination of salt-stressed MTi2 seeds (ANOVA, p<0.05). The optimum KCl concentration is 20-30 mM which has given more than 2x higher percentage and rate of germination, and Seed Vigor compared with the control as presented in Table 4.

Table 4. Salinity-stressed MTi2 seeds treated in KCl: germination percentage, germination rate, and seed vigor

Conc. (mM)	GP (%)	GR	SV
0	37.1°±7.3	6ª±1.4	2.91 ^a ±0.7
10	$51.4^{ab} \pm 10.7$	$8.4^{ab} \pm 1.8$	$4.50^{a}\pm1.2$
20	$74.3^{b} \pm 11.4$	$12.4^{b}\pm1.8$	$10.1^{b}\pm1.8$
30	$82.9^{b}\pm7.0$	$14.2^{b}\pm1.1$	$10.4^{b} \pm 1.0$
40	$65.7^{ab} \pm 5.7$	$10.2^{ab}\pm0.6$	$6.6^{ab}\pm0.5$
50	$62.9^{ab} \pm 7.3$	$9.40^{ab} \pm 1.1$	$5.2^{a}\pm0.7$

The mean is used, and five replicates are used to calculate the standard errors of measurement. Significant differences between the means are shown by superscripts with distinct letters inside the means of each column

(a-c) (Tukey HSD test, p<0.05, Appendix 17-19).

Impact of KCl on the Initial Growth of Salt-Stressed MTi2 Seedlings

Table 5 shows that KCl significantly enhances the growth of salt-stressed MTi2 seedlings (ANOVA, p<0.05). KCl increased significantly in the length of hypocotyls

compared with radicals. The seedling biomass also increased. This study found that the highest seedling length and biomass were found in 20 mM KCl.

Table 5. Salt-stressed MTi2 seedlings in varying KCl concentrations: hypocotyl, radical, seedling length, and biomass

Conc. (mM)	HL (cm)	RL (cm)	SL (cm)	Biomass (g)
0	4.1°±0.4	3.4ª±0.3	7.5°±0.6	0.011 ^a ±0.0021
10	$4.7^{ab}\pm0.7$	$3.9^{ab}\pm0.6$	$8.6^{a}\pm1$	$0.013^{ab} \pm 0.0025$
20	$8.8^{d} \pm 0.6$	$4.6^{ab}\pm0.3$	$13.3^{\circ}\pm0.5$	$0.021^{b}\pm0.001$
30	$7.2^{cd} \pm 0.7$	5.5 ^b ±0.6	$12.7^{bc} \pm 0.9$	$0.018^{ab} {\pm} 0.0032$
40	$6.9^{\text{bcd}} \pm 0.0.5$	$3.3^{a}\pm0.3$	$10.2^{ab} \pm 0.4$	$0.018^{ab} \pm 0.0012$
50	$5.3^{abc} \pm 0.5$	$2.9^{a}\pm0.3$	$8.2^{a}\pm0.3$	$0.013^{ab} \pm 0.0018$

The mean is used, and five replicates are used to calculate the standard errors of measurement. Significant differences between the means are shown by superscripts with distinct letters inside the means of each column (a-c) (Tukey HSD test, p<0.05, Appendix 20-23).

Combination of Salicylic Acid and KCl for the Treatment of Salt-Stressed MTi2 Seeds The percentage and rate of germination, and Seed Vigor

Table 6 shows that the mixture of SA and KCl significantly increased the percentage and rate of germination of salt-stressed MTi2 seeds (ANOVA, p<0.05). The percentage and rate of germination of MTi2 seeds under salt

stress treated with a mixture of SA and KCl increased more than 2x higher than deionized water (control). A mixture of 0.75 mM SA, and 30 mM KCl gave the highest germination percentage and germination rates as presented in Table 6. The mixture of SA and KCl also increased the vigor of salt-stressed MTi2 seeds more than 4x higher than deionized water (control). A combination of 0.50 mM SA and 20 mM KCl gave the highest seed vigor.

Table 6. Salicylic acid and KCl used to treat salt-stressed MTi2 seeds resulted in positive germination percentages, rates, and vigor of seeds

Conc. (mM)	GP (%)	GR	SV
0	37.1°±7.3	$6.0^{a}\pm1.4$	2.91°±0.7
0.50 SA + 20 KCl	$77.1^{b}\pm8.6$	$13.5^{b}\pm1.5$	$15.4^{b}\pm1.8$
0.75 SA + 20 KCl	$80.0^{b}\pm5.7$	$14.1^{b}\pm0.9$	$13.5^{b}\pm1.1$
0.50 SA + 30KCl	82.9 ^b ±8.3	$14.3^{b}\pm1.2$	$13.1^{b}\pm1.6$
0.75 SA +30 KCl	94.3 ^b ±7.8	$16.2^{b}\pm0.4$	13.4 ^b ±0.4

The mean is used, and five replicates are used to calculate the standard errors of measurement. Significant differences between the means are shown by superscripts with distinct letters inside the means of each column (a-c) (Tukey HSD test, p<0.05, Appendix 24-26).

Impact of Salicylic Acid and KCl Combination on Salt Stressed MTi2 Seedlings' Early Seedling Growth:

The combination of SA and KCl increased the early growth of salt-stressed MTi2

seedlings. **Table 7** shows that seedling length is more than 2x higher than control. The mixture also increased the length of hypocotyl, radical, and biomass. The highest seedling growth was found in the mixture of 0.50 mM SA and 20 mM KCl.

Table 7. Salt-stressed MTi2 seedlings in varying SA and KCl concentrations: hypocotyl, radical, seedling length, and biomass

Conc. (mM)	HL (cm)	RL (cm)	SL (cm)	Biomass (g)
0	4.1ª±0.4	3.4°±0.3	7.5°±0.6	0.011°±0.002
0.50 SA + 20 KCl	$13.1^{d}\pm0.3$	$6.8^{\circ} \pm 0.3$	$19.9^{d} \pm 0.4$	$0.024^{\circ} \pm 0.0004$
0.75 SA + 20 KCl	$10.3^{bc} \pm 0.5$	$6.6^{bc} \pm 0.5$	$16.9^{\circ}\pm0.5$	$0.022^{bc}\pm0.0009$
0.50 SA + 30 KCl	$10.7^{\circ} \pm 0.6$	$5.0^{ab}\pm0.3$	$15.7^{bc} \pm 0.6$	$0.020^{bc} \pm 0.001$
0.75 SA + 30 KCl	$8.6^{b}\pm0.5$	$5.7^{bc} \pm 0.5$	$14.3^{b}\pm0.6$	$0.019^{b}\pm0.0005$

The mean is used, and five replicates are used to calculate the standard errors of measurement. Significant differences between the means are shown by superscripts with distinct letters inside the means of each column (a-c) (Tukey HSD test, p<0.05, Appendix 27-30).

Discussion

This experiment was carried out to increase the germination of salt-stressed MTi2 seeds by using SA, KCl, and a mixture of SA and KCl. The application of SA alone and KCl alone increased all germination parameters compared with the control. The ideal concentration for SA is 0.5 mM, and 0.75 mMA SA, while for KCl it is 20 and 30 mM of KCl. Moreover, the mixture of SA and KCl gave higher germination parameters compared with the control and SA alone and KCl alone.

Salicylic acid (SA) is a plant-produced phenolic compound and a plant hormone that acts as the main factor in the induction of plant defense against a multiple of biotic and abiotic stresses through morphological, physiological, and biochemical mechanisms [17, 28, 29]. War [30] reported that SA induced higher activity of peroxidase (POD), polyphenol oxidase (PPO), and amounts of total phenols, H₂O₂, and protein content in chickpea plants. The results suggest that SA at low concentrations could be used for the induction of a plant defensive system that would enable the plant to withstand various biotic and abiotic stresses.

The application of SA under saline conditions caused higher oxygen uptake, greater α -amylase activity, and the efficient mobilization of nutrients from the cotyledons to the embryonic axis and the higher contents of soluble sugar, protein, and free amino acids [31].

High salinity delayed germination but the inhibitory effect was reduced by low concentrations of SA [32]. Under high salinity, osmotic and ionic stresses may induce this stress and then may generate secondary effects such as oxidative stress [33]. SA promotes germination by reducing osmotic damage [32]. Other than that, the SA's exogenous application may induce the metabolic consumption of soluble sugars to form new cell constituents as a mechanism to enhance growth [17].

SA treatments activate a greater accumulation of free amino acids, including proline in the two organs (root and shoot) of the stressed plant. This compound is important as it is involved in osmotic adjustment in the presence of sodium ions. Proline (osmoregulant) is also involved in reducing the injurious effects of salinity and speeding up the repair processes following stress [17]. Proline may also contribute to the development of the antistress reaction [34].

El-Tayeb [17] found that seeds that are exposed to salt will accumulate ion content such as Na+ in the shoot and root. The accumulation of the ions causes a decrease in important nutrients of the plant tissue, for example, K, Ca, and P content. The application of SA reduces the Na+ content, thus indicating SA as the decreasing factor of toxicity absorption. Low content of toxicity elements in plant tissue results in low membrane injury, high water content, and dry matter production. ATP content in the shoot and root of the stressed seed increased, thus it reflects the high content of phosphorus after SA was applied.

Previous studies by Pareek [35]; Hussein [36]; and Mutlu [37] found that applying SA effectively improved the synthesis of soluble proteins, consequently enhancing plant adaptation to stress. Thus, increasing SA to stimulate protein accumulation in salt-stressed plants could lead to better resistance to salinity stress. Studies by El-Tayeb [17]; Deef [38]; and Ashraf [39] have shown that SA plays an influential role in improving crop resistance to salt stress, particularly at the seedling stage Several studies have indicated that the negative impact of salt stress on plants could be reduced by exogenous application of SA [35, 40, 39, 41, 42]. Bahrani [43]; and Baninasab [31] reported that lower doses of SA raised the seed germinates in cucumber, while extremely high concentrations reduced the germination percentage. Similarly, Farahbakhsh [44] in a study of Foeniculum vulgare reported that 0.5 mM of SA in measured traits was more effective than other levels.

Similar results were shown by Naz [45] who studied seed germination and seedling growth of Pisum sativum and found that the adverse effects of salinity were eliminated by applying potassium through seed priming with KCl, which can be a beneficial technique for the activation of germination of maize seeds, leading to safe seedling stands under salt stress during germination, which were enhanced using this priming process [46]. According to Leigh [47]; and Wang [48], the role of potassium is to offer the appropriate ionic environment for metabolic processes in the cytoplasm and therefore act as a regulator of numerous processes such as growth regulation. Both K and Cl are the major ions involved in the neutralization of charges and as the most important inorganic osmotically active ions in plant cells and tissues [49, 50]. K and Cl play an important role in the opening and closing of stomata [51, 52].

K is a plant nutrient that plays a particular role as a surviving mechanism of crop plants under unfavorable stress conditions. The larger K requirement of plants under different abiotic stresses appears to be related to the inhibitory role of K against ROS production. NaCl treatments decreased the K nutrition of plants and suggested that K deficiency at the cellular level might be a contributory factor to salt-induced oxidative stress and related cell damage. Improving the K nutrition of plants under salt stress could be a remarkable strategy for minimizing oxidative cell damage, at least in part by reducing ROS formation [53].

Conclusion

Salinity stress can seriously affect the germination characteristics of MTi2 seeds. Thus, another aim of the current study is to identify an effective therapy that will boost the germination of MTi2 seeds that have been salt-stressed. According to the results, the best concentrations for boosting the germination and early growth of MTi2 seedlings in comparison to the control treatment were found to be 0.5-0.75 mM SA and 20-30 mM KCl. Furthermore, MTi2 seedling germination and early growth were more than 1x higher when the best concentrations of KCl (20-30 mM) and SA (0.5-0.75 mM) were combined. Salicylic acid (SA) and low levels of KCl applied to salt-stressed MTi2 seeds can help reduce the negative effects of salinity stress and enhance the percentage, rate, vigor, length, and biomass of the seedlings that germinate. Where maximum germination percentage (94.3%) was recorded in 0.75mM SA and 30mMKCl, the highest germination rate shows in 0.75 mM SA and 30 mM KCl (14.2), the highest value of seed vigor shows in 0.5 mM SA and 20 mM KCl mM (15.4) and the highest length of seedlings shows in 0.5 mM SA and 20 mM KCl mM (19.9cm). Finally, the highest value of biomass was observed at 0.5 mM SA and 20 mM KCl mM (0.0244g).

Salicylic acid (SA) and a low concentration of KCl applied to salt-stressed MTi2 seeds can contribute to mitigating the negative effects of salinity stress and enhance seed vigor, germination rate, percentage, dry weight, and seedling length.

Conflicts of Interest

The authors hereby declare no conflicts of interest.

Authors Contribution:

All authors contributed in the experiments, analysis and preparation of this manuscript.

Funding/Support

Not Applicable.

References

- Vorasoot N, Songsri P, Akkasaeng C, Jogloy S, and Patanothai A. Effect of water stress on yield and agronomic characters of peanut. Songklanakarin J. Sci. Technol. 2003; 25(3):283-288.
- 2. Kaur G, Kumar S, Nayyar H, and Upadhyaya HD. Cold Stress Injury during the Pod-Filling Phase in Chickpea (Cicer arietinum L.): Effects on Quantitative and Qualitative Components of Seeds. Journal of Agronomy and Crop Science 2008; 194(6):457-464. doi:10.1111/j.1439-037X.2008.00336.x
- 3. Thakur P, Kumar S, Malik JA, Berger JD, and Nayyar H. Cold stress effects on reproductive development in grain crops: an overview. Environmental and Experimental Botany 2010; 67(3):429-443.

doi:10.1016/j.envexpbot.2009.09.004

- 4. Doupis G, Chartzoulakis K, Beis A, and Patakas A. Allometric and biochemical responses of grapevines subjected to drought and enhanced ultraviolet-B radiation. Australian Journal of Grape and Wine Research 2011; 17(1):36-42. doi:10.1111/j.1755-0238.2010.00114.x
- 5. Brinkman R. Saline and sodic soils. In: Land reclamation and water management International Institute for Land Reclamation and Improvement (ILRI). Wageningen, The Netherlands 1980; Pp. 62-68.
- 6. Hakim MA, Juraimi AS, Hanafi MM, Ali E, Ismail MR, Selamat A, and Karim SR. Effect of salt stress on morpho-physiology, vegetative growth and yield of rice. Journal of Environmental Biology 2014; 35(2):317-326. PMID: 24665756

- Flowers TJ, and Muscolo A. Short Communication Special Issue: Physiology and Ecology of Halophytes - Plants Living in Salt-Rich Environments Introduction to the Special Issue: Halophytes in a changing world 2015; 7:1-5. doi: 10.1093/aobpla/plv020.
- 8. Rowell DL, and Wild A. Soil acidity and alkalinity. Russell's soil conditions and plant growth. Eleventh ed. 1988; 844-898.
- Hossain N, Muhibbullah M, Ali KMB, and Molla MH. Relationship between Soil Salinity and Physico-chemical Properties of Paddy Field Soils of Jhilwanja Union, Cox's Bazar, Bangladesh. Journal of Agricultural Science 2015; 7(10):166-180. doi:10.5539/jas.v7n10p166
- 10. Zhao J, Ren W, Zhi D, Wang L, and Xia G. Arabidopsis DREB1A/CBF3 bestowed transgenic tall fescue increased tolerance to drought stress. Plant Cell Reports 2007; 26(9):1521-1528. doi:10.1007/s00299-007-0362-3
- 11. Hassan NA, Drew JV, Knudsen D, and Olson RA. Influence of soil salinity on production of dry matter and uptake and distribution of nutrients in barley and corn: I. Barley (Hordeum vulgare L.). Agronomy Journal 1970; 62(1):43-45.
- 12. Sato S, Sakaguchi S, Furukawa H, and Ikeda H. Effects of NaCl application to hydroponic nutrient solution on fruit characteristics of tomato (Lycopersicon esculentum Mill.). Scientia horticulturae 2006; 109(3):248-253. doi:10.1016/j.scienta.2006.05.003
- 13. Yalpani N, Enyedi AJ, León J, and Raskin I. Ultraviolet light and ozone stimulate accumulation of salicylic acid, pathogenesis-related proteins and virus resistance in tobacco. Planta 1994; 193(3):372-376.
- 14. Matsuoka M. Gibberellin signaling: how do plant cells respond to GA signals. Journal of Plant Growth Regulation 2003; 22(2):123-125. doi:10.1007/s00344-003-0039-2
- 15. Shakirova FM, Sakhabutdinova AR, Bezrukova MV, Fatkhutdinova RA, and Fatkhutdinova DR. Changes in the hormonal status of wheat seedlings induced by salicylic acid and salinity. Plant Science

- 2003; 164(3):317-322. doi:10.1016/S0168-9452(02)00415-6
- 16. Srivastava NK, and Srivastava AK. Influence of gibberellic acid on 14CO2 metabolism, growth, and production of alkaloids in Catharanthusroseus. Photosynthetica 2007; 45(1):156-160. doi:10.1007/s11099-007-0026-0
- El-Tayeb MA. Response of barley grains to the interactive e.ect of salinity and salicylic acid. Plant Growth Regulation 2005; 45(3):215-224. doi:10.1007/s10725-005-4928-1
- 18. Szepesi Á, Csiszár J, Gémes K, Horváth E, Horváth F, Simon ML, and Tari I. Salicylic acid improves acclimation to salt stress by stimulating abscisic aldehyde oxidase activity and abscisic acid accumulation, and increases Na+ content in leaves without toxicity symptoms in Solanum lycopersicum L. Journal of plant physiology 2009; 166(9):914-925.

doi:10.1016/j.jplph.2008.11.012

- 19. Grubben GJ, Denton OA. Plant resources of tropical Africa 2. Vegetables. Plant resources of tropical Africa 2. Vegetables 2004; 667.
- 20. Panuccio MR, Jacobsen SE, Akhtar SS, and Muscolo A. Effect of saline water on seed germination and early seedling growth of the halophyte quinoa. Annals of Botany 2014; 6:plu047.

doi:doi.org/10.1093/aobpla/plu047

- 21. Mohammed SJ, and Nulit R. Seed Priming Improves the Germination and Early Growth of Turnip Seedlings under Salinity Stress Periódico Tchê Química 2020; 17(35):73-82. doi:10.52571/PTQ.v17.n35.2020.07_SAMAR_pgs _73_82.pdf
- 22. Afzal I, Rauf S, Basra SMA, and Murtaza G. Halopriming improves vigor, metabolism of reserves and ionic contents in wheat seedlings under salt stress. Plant Soil Environ 2008; 54(9):382-388.
- 23. Elouaer MA, and Cherif H. Effect of NaCl priming duration and concentration on germination behavior of Tunisian safflower. Journal of Stress Physiology & Biochemistry 2012; 8(3): 2.

- 24. Achakzai AKK. Effect of water stress on imbibition, germination and seedling growth of maize cultivars. Sarhad J. Agric 2009; 25(2):165-172.
- 25. Kandil AA, Sharief AE, Abido WAE, and Ibrahim MM. Effect of salinity on seed germination and seedling characters of some forage sorghum cultivars. International Journal of Agriculture Sciences 2012; 4(7):306-311. ISSN: 0975-3710 & E-ISSN: 0975-9107
- 26. Dezfuli P M, Sharif-Zadeh F, and Janmohammadi M. Influence of priming techniques on seed germination behavior of maize inbred lines (Zea mays L.). Journal of Agricultural and Biological Science 2008; 3(3):22-25.
- 27. Abdul-Baki AA, and Anderson JD. Vigor determination in soybean seed by multiple criteria. Crop Science 1973; 13(6):630-633.
- 28. Hayat S, Ali B, and Ahmad A. Salicylic acid: biosynthesis, metabolism and physiological role in plants. In Salicylic acid: A plant hormone. Springer, Netherlands 2007; pp. 1-14. doi:10.1007/1-4020-5184-0 1
- 29. Gharib FA, and Hegazi AZ. Salicylic acid ameliorates germination, seedling growth, phytohormone and enzymes activity in bean (Phaseolus vulgaris L.) under cold stress. Journal of American Science 2010; 6(10):675-683.
- 30. War AR, Paulraj MG, War MY, and Ignacimuthu S. Role of salicylic acid in induction of plant defense system in chickpea (Cicer arietinum L.). Plant signaling & behavior 2011; 6(11):1787-1792. doi:10.4161/psb.6.11.17685
- 31. Baninasab B, and Baghbanha MR. Influence of salicylic acid pre-treatment on emergence and early seedling growth of cucumber (Cucumis sativus) under salt stress. International Journal of Plant Production 2013; 7(2).
- 32. Lee S, Kim SG, and Park CM. Salicylic acid promotes seed germination under high salinity by modulating antioxidant activity in Arabidopsis. New Phytologist 2010; 188(2):626-637. doi:10.1111/j.1469-8137.2010.03378.x

- 33. Zhu JK. Genetic analysis of plant salt tolerance using Arabidopsis. Plant Physiology 2000; 124(3):941-948. doi:10.1104/pp.124.3.941
- 34. Hara M, Furukawa J, Sato A, Mizoguchi T, and Miura K. Abiotic stress and role of salicylic acid in plants. In Abiotic Stress Responses in Plants. Springer, New York 2012; pp. 235-251. doi:10.1007/978-1-4614-0634-1_13
- 35. Pareek A, Singla, SL, and Grover A. Salt Responsive Proteins/Genes In Crop Plants.In Strategies For Improving Salt Tolerance In Higher Plants (eds P.K.) Jaiwal, R. P. singh, and A. Gulati, Oxford and IBH Puplication Co.,New Delhi 1997; 365-391.
- 36. Hussein MM, Balbaa LK, and Gaballah MS. Salicylic acid and salinity effects on growth of maize plants. Research Journal of Agriculture and biological Sciences 2007; 3(4), 321-328.
- 37. Mutlu S, Karadağoğlu Ö, Atici Ö, and Nalbantoğlu B. Protective role of salicylic acid applied before cold stress on antioxidative system and protein patterns in barley apoplast. Biologia Plantarum 2013; 57(3):507-513. doi:10.1007/s10535-013-0322-4
- 38. Deef HE. Influence of salicylic acid on stress tolerance during seed germination of Triticum aestivum and Hordeum vulgare. Advances in biological research 2007; 1 (1-2):40-48.
- 39. Ashraf M. Registration of 'S-24' spring wheat with improved salt tolerance. Journal of plant registrations 2010; 4(1):34-37. doi:10.3198/jpr2008.05.0252crc
- 40. Noreen S, and Ashraf M. Alleviation of adverse effects of salt stress on sunflower (Helianthus annuus L.) by exogenous application of salicylic acid: growth and photosynthesis. Pakistan Jurnal Botany 2008; 40(4):1657-1663.
- 41. Pirasteh-Anosheh H, and Emam Y. Manipulation of morpho-physiological traits in bread and durum wheat by using PGRs at different water regimes. J. Crop Prod. Process 2012; 5:29-45. http://dorl.net/dor/20.1001.1.22518517.1391.2.5.3.1
- 42. Pirasteh-Anosheh H, Emam Y, Ashraf M, and Foolad MR. Exogenous application of salicylic acid

- and chlormequat chloride alleviates negative effects of drought stress in wheat. Adv. Stud. Biol. 2012; 11(4):501-520.
- 43. Bahrani A, and Pourreza J. Gibberlic acid and salicylic acid effects on seed germination and seedlings growth of wheat (Triticum aestivum L.) under salt stress condition. World App Sci J 2012; 18(5):633-641.
- 44. Farahbakhsh H. Germination and seedling growth in un-primed and primed seeds of fennel as affected by reduced water potential induced by NaCl. International Research Journal of Applied and Basic Sciences 2012; 3(4): 737-744.
- 45. Naz F, Gul H, and Hamayun M. Effect of NaCl Stress on Pisum sativum Germination and Seedling Growth with the Influence of Seed Priming with Potassium (KCl and KOH). American-Eurasian Journal Agriculture & Environ Science 2014;14 (11):1304-1311. doi:10.5829/idosi.aejaes.2014.14.11.748
- 46. Ali A, Hyder SI, Arshadullah M, and Bhatti SU. Potasssium chloride as a nutrient seed primer to enhance salttolerance in maize. Pesquisa Agropecuária Brasileira 2012; 47(8):1181-1184. doi:10.1590/S0100-204X2012000800020
- 47. Leigh RA, and Wyn-Jones RG. A hypothesis relating critical potassium concentrations for growth to the distribution and functions of this ion in the plant cell. New Phytologist 1984; 97(1):1-13.
- 48. Wang Y, and Wu WH. Potassium transport and signalling in higher plants. Annual Review of Plant Biology 2013; 64:451-476. doi:10.1146/annurev-arplant-050312-120153
- 49. Clarkson DT, and Hanson JB. The mineral nutrition of higher plants. Annual review of plant physiology 1980; 31(1):239-298.
- 50. Mitra GN. Chloride (Cl-) Uptake. In Regulation of Nutrient Uptake by Plants, Springer India 2015; Pp. 167-173. doi:10.1007/978-81-322-2334-4_17
- 51. Talbott LD, and Zeiger E. Central roles for potassium and sucrose in guard-cell osmoregulation. Plant Physiology 1996; 111(4):1051-1057.

- 52. Santelia D, and Lawson T. Rethinking Guard Cell Metabolism. Plant Physiology 2016; 172(3):1371-1392. doi:10.1104/pp.16.00767
- 53. Cakmak I. The role of potassium in alleviating detrimental effects of abiotic stresses in plants. Journal of Plant Nutrition and Soil Science 2005; 168(4):521-530. doi:10.1002/jpln.200420485