

## Phyto-pharmacological Benefit of Methanol Whole Plant Extract of *Ocimum canum* Sims. in Chemically-induced Epileptic Rat Model

Yakubu James<sup>1</sup>, Mbaya Abdullahi Hassan<sup>2</sup>, Yesufu Hassan Braimah<sup>3</sup>,  
Abdulrahman Fanna Inna<sup>4</sup>, Sodipo Olufunke Adebola<sup>5</sup>, Medugu Asinamai Ndai<sup>6</sup>,  
Wampana Bitrus<sup>7</sup>

<sup>1</sup>Department of Pure and Applied Chemistry, Faculty of Science, University of Maiduguri, Borno State, Nigeria. E-mail: jamesyakubu96@gmail.com

<sup>2</sup>Department of Pure and Applied Chemistry, Faculty of Science, University of Maiduguri, Borno State, Nigeria. Email: abdullahihassan770@gmail.com

<sup>3</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Maiduguri, Maiduguri, Borno State, Nigeria. E-mail: hassanyesufubraimah@gmail.com

<sup>4</sup>Department of Pure and Applied Chemistry, Faculty of Science, University of Maiduguri, Borno State, Nigeria. Email: fannas06@gmail.com

<sup>5</sup>Department of Clinical Pharmacology and Therapeutics, Faculty of Basic Clinical Sciences, College of Medical Sciences, University of Maiduguri, Borno State, Nigeria. E-mail: sodipoolufunke@yahoo.com

<sup>6</sup>Department of Pharmaceutical Outcomes and Policy, College of Pharmacy, University of Florida, Florida, USA. E-mail: a.ndai@ufl.edu

<sup>7</sup>Department of Pharmacology, Faculty of Veterinary Medicine, University of Maiduguri, Nigeria. E-mail: bwampana@gmail.com  
Corresponding Author: Department of Pure and Applied Chemistry, Faculty of Science, University of Maiduguri, Borno State, Nigeria. E-mail: jamesyakubu96@gmail.com

### Article Info

#### Article type:

Original Article

#### Article History:

**Received:** 07 Jan 2023

**Received in revised form:**

17 March 2023

**Accepted:** 28 April 2023

**Published online:** 31 May

2023

### Abstract

**Objectives:** The role of *Ocimum canum* Sim., as a medicinal agent used for the treatment and management of diseases cannot be overemphasized. The aim of the study is to screen the Phytochemical constituents and to evaluate the antiepileptic effect of methanol whole plant extract in Wistar albino rats.

**Material and Methods:** The whole plant part of *O. canum* was extracted with 85% percentage of methanol and screened for phytochemicals. An acute toxicity study was carried out using Lorke's method and the antiepileptic activity was evaluated using pentylenetetrazole (PTZ) and strychnine in rats. One-way analysis of variance (ANOVA) was used in the analysis of the data.

**Results:** The phytochemical analysis revealed the presence of saponins, cardiac glycosides, flavonoids, terpenoids, and cardenolides. The intraperitoneal median lethal dose value (LD50) of *O. canum* in rats was 2154 mg/kg. The extract at the dose of 300 and 600 mg/kg body weight protected 40% and 80% of experimental animals against both PTZ and strychnine-induced convulsion; protected 60% and 80% of rats



**Keywords:** against death induced by strychnine; compared to 100% protection by the standard drug sodium valproate (20mg/kg).  
*Ocimum canum*,  
 Phytochemical,  
 Antiepileptic,  
 Pentylenetetrazole (PTZ),  
 Strychnine

**Conclusion:** The antiepileptic investigation suggests that methanol extract of whole plant of *O. canum* has antiepileptic activity which could be employed for the management of epileptic related conditions.

## Introduction

Plants have had been a source of food, shelter and medicine. The use of plants medicinally was mostly developed through observing wild animals and trying and failing [1].

The surge in the use of plant in medication formulation is due to the availability, affordability, accessibility, safety, efficiency, and rare side effects of drugs from plants [2]. A lot of drugs listed as conventional medications were derived from plants originally [3]. In Africa, up to 80 % of the population uses traditional medicine for primary health care. Traditional medicine has maintained its popularity in all regions. In Nigeria, Ghana, Mali, and Zambia for example, the first line of treatment for 60 % of children with high fever resulting from malaria, epilepsy and cold is the use of herbal medicines at home [1].

Epilepsy is a major neurological disorder and up to 5% of the world population develop epilepsy in their lifetime [5]. It is a disease that affects about 50 million

people across the globe and 85% of this population resides in developing countries, it is the second commonest neurological disorder. The World Health Organization [6] has estimated that 5 million people are diagnosed with epilepsy each year. In Nigeria, the estimated prevalence of epilepsy is 8 per 1000 people indicating a substantial burden of the disease in the country [7]. Available antiepileptic drugs (AED) are synthetic chemicals that have overwhelming side effects such as weight gain, hepatotoxicity, teratogenicity, and withdrawal symptoms [8]. The treatment of epilepsy with available AED is symptomatic as these drugs only prevent seizure and do not cure the underlying disease process in the brain [9]. It is generally estimated that up to 30% of patients are refractory to conventional AED treatment [10]. Although a broad range of newer and more selective agents are currently being used, there is still a need for more selective and less toxic AEDs. Several plants used for the treatment of epilepsy in different systems of traditional medicine have shown activity when tested on modern bioassays for the detection of anticonvulsant activity [4] and many such plants remain to be scientifically investigated.

*O. canum* Sims. (Figure 1) belongs to the family. It is a small herbaceous, erect, perennial, and aromatic plant usually growing 20 – 30 cm tall with some specimens up to 100 cm [11]. The plant is often found in tropical climates, growing on roadsides, in fields, in teak forests, and in open waste place close to settlements. It also often grown as an annual, especially



**Figure 1:** *Ocimum canum* Sims in its natural habitat

in cooler climates. The entire plant is highly aromatic, with an odour comparable to citrus. As such, it can be used for culinary purposes in similar ways to sweet basil (*O. basilicum*) [11]. The aromatic leaves are used

The leaves can also be used as a flavouring in salads, sauces, soups etc. The young leaves, which are eaten raw as a vegetable side-dish in Indonesia, Malaysia, and Thailand. Due to the high fragrance nature of the leaves, it is also added to various dishes with fishy or unpleasant smell. Medicinally, the leaves of the plant are used for several ailments. The decoctions are used to treat coughs and for respiratory related conditions, the whole plant is used in baths to treat rheumatism, renal colic, and calcification [12]. Pharmacological investigation which include anaesthetic [13], antimicrobial [14], antioxidant [15], anti-inflammatory [16], anxiolytic [17], and insecticidal effect [18] of the plant has been carried out. This study aimed to scientifically justify the use of this plant for the acclaimed use for the treatment and management of convulsion and related disease in Northeastern Nigeria.

## Materials and Methods

### Experimental Site

The chemical and pharmacological experiments were conducted in the laboratories of the Department of Pure and Chemistry and Faculty of Pharmacy, University of Maiduguri, Borno State, Nigeria respectively.

### Plant Collection and Identification and Preparation

The fresh whole plant *O. canum* was collected from Maiduguri from Konduga Local Government Area, Borno State, Nigeria. The identity of the plant was verified by Professor S.S. Sanusi; a Taxonomist of the

as a flavouring in a range of foods, whilst a cooling drink can be made from the seeds. The plant is gathered from the wild and also often cultivated [11].

Department of Biological Sciences, University of Maiduguri, Borno State, Nigeria. A voucher specimen was deposited at the Herbarium of the Department of Pharmacognosy, Faculty of Pharmacy, University of Maiduguri for future reference. The sample was air-dried under shade for a period of 2 weeks and was size-reduced into coarse powder using mortar and pestle.

### Extraction and Preliminary Phytochemical Screening

Two hundred and fifty grammes (250g) of the powdered whole plant was soxhlet extracted using 95% methanol for 6hrs. The obtained solution was then filtered to remove debris, and was concentrated under reduced pressure and the yield was 31g. The extract was transfer into an air-tight container until required for analysis.

The extract was subjected to preliminary phytochemical screening to test for the presence of secondary metabolites, which include: alkaloids, carbohydrates, flavonoids, saponins, tannins, glycosides, (cardiac, steroidal), and terpenes/terpenoids using conventional protocols [19,20,21].

### Anticonvulsant Study

#### Effect of Methanol Whole Plant Extract of *Ocimum canum* on Pentylene-tetrazole-Induced Convulsion in Rats

Experimental method of Anticonvulsant study as described by Swinyard *et al.* [22]; Medugu *et al.* [22] and Yakubu *et al.* [24] was adopted in this study. Twenty-five (25) albino rats of both sexes weighing between 100-220 g were used. They were housed in a

clean cage and were given food and water *ad libitum*. They were then divided into five groups of five rats and labelled as group A, B, C, D, and E. Group A was given 100 mg/kg of pentylenetetrazole (PTZ) *i.p* and served as the negative control and were not pre-treated. Group B, C, and D were pre-treated intraperitoneally with 300 mg/kg and 600 mg/kg of the extract respectively, 30 minutes before treatment with convulsant (100 mg/kg of PTZ) and Group F was given 5 mg/kg orally of sodium valproate which serve as the positive control as standard anticonvulsant drug. During the experiment the onset of convulsion, number of convulsions per minute, and the duration of convulsions were recorded for 30minutes. Also the number of animals which survived within the period of observation were expressed as percentage (%) protection.

### Effect of Methanol Whole Plant Extract of *Ocimum canum* on Strychnine-Induced Convulsion in Rats

The method described by Medugu *et al.* [23] was adopted in this study. In brief, strychnine convulsion was induced by the subcutaneous injection of 1mg/kg of strychnine nitrate in the rats. 30min prior to administration, 2 groups (groups 3 and 4) of 5 animals each were intraperitoneally pre-treated with methanol extract of *Ocimum canum* with doses of 300mg/kg and 600mg/kg. The fourth group was treated with sodium valproate (20mg/kg *i.p*) which served as the positive control while the fifth group received normal saline 10ml/kg as the negative control. The rats were observed for tonic extensor jerks of the hind limbs followed by death in 30 minutes. Abolition of tonic extensor jerks of the hind limb was considered an indicator that the extract could prevent strychnine-induced convulsion.

### Data Handling

The generated data from the PTZ and strychnine-induced antiepileptic study were express as Mean  $\pm$  Standard Error of the mean and analyzed by one-way analysis of variance (ANOVA) using Statistical Graphpad Prism Version 9.0 for Windows. Probability value of  $P < 0.05$  were considered significant.

## Results and Discussion

### Phytochemical Screening of Methanol

#### Whole Plant Extract of *Ocimum canum*

The phytochemical screening of the methanol extract obtained from the whole plant of *Ocimum canum* revealed the presence of phytochemicals which include flavonoids, terpenoids, cardiac glycosides, cardenolides, carbohydrates, and saponins. The result is presented in Table 2.

### Acute Toxicity of Methanol Whole Plant Extract of *Ocimum canum*

The result for acute toxicity ( $LD_{50}$ ) study obtained is shown in Table 3. The oral and intraperitoneal  $LD_{50}$  were  $\geq 5000$  mg/kg, as no death was recorded and 3807mg/kg as death occurred at 5000mg/kg dose of the extract respectively.

### The Effect of Methanol Leaf Extract of *Ocimum canum* on PTZ-Induced Convulsion in Rats

The methanol extract significantly conferred protection of 40% and 80% against PTZ-Induced convulsion when treated with extract doses of 300 mg/kg and 600 mg/kg respectively in a dose dependent manner. There was also a decrease in the number of convulsion, onset of convulsion while there was a significant increase in time of death. The result of this study is presented in Table 3. The Effect of Methanol

Extract of *O. canum* on Strychnine-Induced Convulsion in Rats

The extract of *O. canum* at doses of 300 and 600mg/kg conferred 40% and 80% respectively against strychnine-induced death. The extract acted by

significantly ( $P < 0.05$ ) prolonging the onset of convulsion and time of death in a dose-dependent manner; while sodium valproate (+ve control drug) had 100% protection effect against strychnine induced convulsion. Result of the study is shown in Table 4.

**Table 1:** *Phytochemical Analysis of O. canum Whole Plant Methanol Extract*

S/N	Phytochemical Test	Inference
1	Carbohydrates	+
2	Tannins	–
3	Phlobatannins	–
4	Cardiac glycosides	+
5	Flavonoids	+
6	Terpenes	+
8	Saponins	+
9	Soluble starch	–
10	Alkaloids	–
11	Steroidal nucleus	+

**Table 2:** *Intraperitoneal (IP) Acute Toxicity Effect of O. canum whole Plant Methanol Extract*

Phase	Dose (mg/kg)	No. of rat	Mortality rate
			IP route
<b>I</b>	10	3	0/3
	100	3	0/3
	1000	3	1/3
<b>II</b>	600	1	0/1
	1000	1	0/1
	16000	1	0/1
	2900	1	1/1

*Ip* LD<sub>50</sub> = 2154 mg/k

**Table 3:** Effect of *O. canum* Whole Plant Extract of Methanol on Pentylene-tetrazole-induced Epilepsy in Rats

Extract pretreatment (mg/kg)	Mean±SEM onset of spasm (min)	Mean±SEM onset of convulsion (min)	Mean±SEM onset of death (min)	Quantal death	Survival (%)
Control Vehicle	7.94±0.32	7.48±1.00	18.20±0.46	1/5	20
Control + 100 mg/kg of PTZ	18.54±0.47	7.32±1.00	17.58±0.25	1/5	20
300 mg/kg + 100 mg/kg of PTZ	11.68±2.07	12.18±0.30	21.34±0.35	2/5	40
600 mg/kg + 100 mg/kg of PTZ	17.68±0.34	18.24±0.41	36.82±10.02	4/5	80
Sodium Valproate (20mg/kg)	7.94±0.32	0.20±0.20	0.00±0.00	5/5	100

Results are in mean±standard error of mean (SEM); n=5; data with the same superscript across columns are statistically not significant (p>0.05)

**Table 3:** Effect of *O. canum* Whole Plant Methanol Extract on Strychnine-Induced Epilepsy in Rats

Extract pretreatment (mg/kg)	Mean±SEM onset of spasm (min)	Mean±SEM onset of convulsion (min)	Mean±SEM onset of death (min)	Quantal death	Survival (%)
Control Vehicle	7.94±0.32	7.48±1.00	18.20±0.46	1/5	20
Control + 100 mg/kg of PTZ	18.54±0.47	7.32±1.00	17.58±0.25	1/5	20
300 mg/kg + 100 mg/kg of PTZ	11.24±0.60	15.20±0.37	24.50±0.21	2/5	40
600 mg/kg + 100 mg/kg of PTZ	6.36±0.16	21.70±70	27.02±0.39	3/5	80
Sodium Valproate (20mg/kg)	7.94±0.32	0.20±0.20	0.00±0.00	5/5	100

Results are in mean±standard error of mean (SEM); n=5; data of same column are all statistically significant (p<0.05)



The present phytochemical study of the methanol whole plant *O. canum* showed that the plant contains phytochemicals of pharmacological importance and has garnered scientific interest, these includes flavonoids, cardiac glycosides, tannins, saponins, and steroids. This report is supported by the findings of [25] whose study revealed the presence of coumarins, flavanoids, glycosides, phenols, tannins, saponins, and steroids, while alkaloids were not detected. Therapeutic benefits of traditional remedies might depend upon individual or combination of these constituents [26], such as those identified in the extract of this plant.

The most abundant phytochemical detected in the present work is terpenoids. It is one of the most abundant phytochemical found in mostly leaves and seeds of plants as volatile oils. It acts as phyto-alexins in plant direct defense responses which involves herbivores and their natural enemies [27]. They are reported to have neuro-pharmacological activity including anticonvulsant activity [28]. Flavonoids is another interesting class of plants' chemical having broad biological and pharmacological activities which act as powerful antioxidants which helps to protect the human body from free radicals and reactive oxygen species [29]. They have been found to inhibit almost all the mechanisms involved in seizures generation in epilepsy. They have been found to modulate neuronal Na<sup>+</sup> channels [30], Ca<sup>2+</sup> channels [31], GABAergic pathway glutamatergic pathway and opioid pathway [32].

The intraperitoneal median lethal dose value (LD<sub>50</sub>) of *O. canum* in mice was 2154 mg/kg. According to Clarke and Clarke [33] any substance whose LD<sub>50</sub> is greater than 1,000 mg/kg is considered non-toxic and can be said to be a good candidate for further research or as a drug or food.

Protection of the experimental animals against pentylenetetrazole (PTZ)-induced seizure predicts anticonvulsant activity and the delayed onset of seizure indicates that it can raise seizure threshold [34]. The anticonvulsant activity of *O. canum* at doses of 300 mg/kg and 600 mg/kg intraperitoneally (i.p) was studied by PTZ-induced seizure model. Pentylenetetrazole (PTZ) is an antagonist of gamma-aminobutyric acid receptors that induces acute convulsions in rodents [35] and it also reduces the T-type of Ca<sup>++</sup> currents [36]. Since the methanol whole plant extract of *O. canum* delayed the occurrence of PTZ-

induced convulsion, it is likely that it may be interfering with GABAergic mechanism to exert its anticonvulsant activity.

Strychnine is a known convulsant which directly antagonizes the inhibitory spinal reflexes of glycine [37]. The crude extract of the whole plant of *O. canum* at doses of 300 mg/kg and 600 mg/kg protected 60% and 80% of the rats against strychnine-induced death. Strychnine induces convulsion by interfering with postsynaptic inhibition mediated by glycine which is an important inhibitory transmitter of the motor neurons and interneurons in the spinal cord. It acts as a competitive and selective antagonist at all glycine receptors [38]. The ability of the extract to prevent the strychnine-induced seizures amasses the anticonvulsant effects mediated via glycine receptors [39].

## Conclusion

The methanol extract of the whole plant of *O. canum* contains saponins, cardiac glycosides, flavonoids, terpenoids, and cardenolides. The whole plant methanol extract had no observable toxic effect on rats within the duration of time evaluated. The anticonvulsant property of the methanol extract was evaluated using two animal models – Pentylenetetrazole and strychnine. The study revealed a significant and promising anticonvulsant effect of the extract which could be attributed to the presence of the individual or synergistic effect of the phytochemicals.

## Conflict of Interest

The authors declared no conflicts of interest.

## Acknowledgement

The authors acknowledge the technical assistance of Mr. Fine Akawu from the Postgraduate Research laboratory, Department of Pure and Applied Chemistry, University of Maiduguri, Maiduguri, Borno State.

## Funding

The research was self-sponsored by the authors. No fund was received from any Organization or Institution for this study.

## References

1. Akinyemi O, Oyewole SO, Jimoh KA. Medicinal plants and sustainable human health: A Review. Horticult Int J 2018; (4):194-195.<https://doi.org/10.15406/hij.2018.02.00051>
2. Dewick PM. Tumor Inhibition from Plants: Trease and Evans Pharmacognosy, 11th Edition, 1996; pp. 45-50.
3. Lucy H, Edgar JD. Medicinal Plants: a re-emerging health aid. Elect J Biotech. 1999; 2(2): 7-14.<https://doi.org/10.2225/vol2-issue2-fulltext-2>
4. Raza M, Shaheen F, Choudhary MI, Sombati, S, Rafiq, A, Suria, A, Rahman A, DeLorenzo RJ. Anticonvulsant activities of ethanolic extract and aqueous fraction isolated from *Delphinium denudatum*. J Ethnopharmacol. 2001; 78: 73 - 78.[https://doi.org/10.1016/S0378-8741\(01\)00327-0](https://doi.org/10.1016/S0378-8741(01)00327-0)
5. Schachter, SC. Evaluation and management of the first Seizure in adults. Up to date. <http://www.update.com/contents/search>. 2021.
6. WHO. World Health Organization. Epilepsy.<https://who.int/news-room/fact-sheets/detail/epilepsy#> 2020.
7. Owolabi, LF, Adamu B, Jibo, AM, Owolabi, DS, Isa, AI, Alhaji, ID, et al. Prevalence of active epilepsy, lifetime epilepsy prevalence, and burden of epilepsy in sub-Saharan Africa from meta-analysis of door-to-door population-based surveys. Epilepsy Behav 2020;103(Pt A):106846.<https://doi.org/10.1016/j.yebeh.2019.106846>
8. McNamara DJ. Pharmacotherapy of epilepsies. In: (Brunton, L.L., Lazo, S.J. and Parker, K.L (Eds) Goodman and Gilman's. The pharmacological basis of therapeutics, Eleventh edition. McGraw-Hill Medical Publishing Division, New York 2006; 501-525.
9. Schmidt D. The clinical impact of new antiepileptic drugs after a decade of use in epilepsy research, 2002; 50: 21-32.[https://doi.org/10.1016/S0920-1211\(02\)00065-7](https://doi.org/10.1016/S0920-1211(02)00065-7).
10. Arroyo S, Brodie MJ, Aranzini G. Is refractory epilepsy preventable? Epilepsia 2002;43:437-444.<https://doi.org/10.1046/j.1528-1157.2002.38501.x>
11. Missouri Botanical Garden. Flora of China Missouri Botanical Garden Press; St. Louis <http://flora.huh.harvard.edu/china/> 1994.
12. Ngassoum, MB, Ousmaila H, Ngamo, LT, Maponmetsem, PM, Jirovetz, L, Buchber G. Aroma compounds of essential oils of two varieties of the spice plant *Ocimum canum* Sims. from northern Cameroon J. Food Comp. Anal 2004; 197-204.<https://doi.org/10.1016/j.jfca.2003.08.002>.
13. de Lima Silva L, Garlet QI, Koakoski G, de Abreu MS, Mallmann CA, Baldisserotto B, et al. Anesthetic activity of the essential oil of *Ocimum americanum* in *Rhamdia quelen* (Quoy & Gaimard, 1824) and its effects on stress parameters. Neotrop Ichthyol 2015; 13(4):715-722.<https://doi.org/10.1590/1982-0224-20150012>.
14. Usman LA, Yusuf AO, Saliu BK, Olanipekun BE. Effect of collection time on chemical composition and antibacterial activity of flower essential oil of *Ocimum canum* (Sims) grown in Nigeria. J Turk Chem Soc Chem 2017; 4(1):149-164.<https://doi.org/10.18596/jotcsa.287322>.
15. Karau GM, Njagi ENM, Machochi AK, Wangai LN, Nthinga, MJ. Chemical composition and in vitro antioxidant activities of *Ocimum americanum*, Adv Analyt Chem 2015; 5(2): 42-49.<https://doi.org/10.5923/j.aac.20150502.03>.
16. Yamada AN, Grespan R, Yamada AT, Silva EL, Saulo Euclides Silva-Filho, Marcio José D, et al. Anti-inflammatory Activity of *Ocimum americanum* L. Essential Oil in Experimental Model of Zymosan-Induced Arthritis. Am J Chinese Med 2013; 41(4): 913-926.<https://doi.org/10.1142/S0192415X13500614>.
17. Barchuk O, Pryshlyak A, Shanaida M. Chemical compositions and sedative activities of the *Dracocephalum moldavica* L. and *Ocimum americanum* L. essential oils, Pharmacologyonline, 2021;2:179-187.
18. Shadia E, El-Aziz A, Omer EA, Sabra SA. Chemical composition of *Ocimum americanum* essential oil and its biological effects against, *Agrotis ipsilon*, (Lepidoptera: Noctuidae). Res J Agri Bio Sci 2007;3(6):740-747.
19. Sofowora EA. Medicinal Plants and Traditional Medicine in Africa. 3rd Edition. Spectrum Books Limited, Ibadan Nigeria 2008; 97-112.
20. Evans WE. Trease and Evans Pharmacognosy. 16th Edition. Saunders Publishers, London 2009; 42-229.



21. Yakubu J, Usman YB, Balami MV, Jonathan, N, Semiu, Y, Teri, DM. Phytochemical and elemental analysis of some organic solvent leaf extracts of *Piliostigma thonningii* (Schumach) Milne-Redhead Ewemen J Folklore Med 2016;2(2): 49-57.
22. Swinyard EA, Woodhead JH, White HS, Franklin, MR. General principles: Experimental selection, quantification and evaluation of anticonvulsant, 3rd ed. In: Levy, RH, Mattson B, Melrum JK, Dreifuss FE. (eds), Springer-Verlag Berlin Heidelberg: New York, USA 1989; 566-874.
23. Medugu AN, J. Yakubu J, Balami VM, Medugu NM, Tata FY, Marte HI et al. Anti-epileptic studies of *Chamaecrista mimosoides* ethanol extract. Tropical J. Nat. Prod. Res 2020;4(5); 203 - 209.<https://doi.org/10.26538/tjnpr/v4i5.4>.
24. Yakubu J, Sodipo, OA, Abdulrahman FI, Balami VM. Toxicity study and anticonvulsant effect of ethanol leaf extract of *Piliostigma thonningii* Milne-Redhead (Fabaceae). Nig J Pharmaceut Res 2021; 17(1); 63-70.<https://doi.org/10.4314/njpr.v17i1.7>
25. Ononamadu CJ, Alhassan AJ, Imam AA, Ibrahim A, Ihegboro GO, Owolarafe AT, et al. In vitro and in vivo anti-diabetic and anti-oxidant activities of methanolic leaf extracts of *Ocimum canum*, Casp J Int Med 2019;10(2):162-175.<https://doi.org/10.22088/cjim.10.2.162>.
26. Amos S, Akah, P, Enwerem N, Chindo B, Hussaini I, Wambebe C, et al. Behavioral effect of *Pavetta crassipes* extract on rodents. Pharmacol Biochem Behav 2004;77:751-759.<https://doi.org/10.1016/j.pbb.2004.01.020>
27. McCaskill D, Croteau R. Prospects for the bioengineering of isoprenoid biosynthesis. Adv Biochem Eng Biotechnol 1997; 55:107-146.<https://doi.org/10.1007/BFb0102064>.
28. Mamta S, Jyoti S, Rajeev N, Dharmendra S, Abhishek G. Phytochemistry of Medicinal Plants. J Pharmacog Phytochem 2013;1(6):168-182.
29. Tapas AR, Sakarkar DM, Kakde RB. Flavonoids and Nutraceuticals: A Review. Trop J Pharmaceut Res 2008;7:1089-1099.<https://doi.org/10.4314/tjpr.v7i3.14693>.
30. Nicholson RA, David LS, Pan RL, Liu XM. Pinostrobin from *Cajanus cajan* (L.) Millsp. inhibits sodium channel-activated depolarization of mouse brain synaptoneurosome. J Fitoterapia 2010; 81:826-9.<https://doi.org/10.1016/j.fitote.2010.05.005>.
31. Cogolludo A, Frazziano G, Briones AM, Cobeno L, Moreno L, Federica L. The dietary flavonoid quercetin activates BKCa currents in coronary arteries via production of H<sub>2</sub>O<sub>2</sub>; Role in vasodilatation. J Cardiovasc Res 2007; 73:424-31.<https://doi.org/10.1016/j.cardiores.2006.09.008>.
32. Engelborghs S, Hooge RD, Deyn PPD. Pathophysiology of epilepsy. J Acta Neurol Belg 2000; 100: 201-213.
33. Clarke, ECG, Clarke, MI. Veterinary Toxicology 2nd edition. Bellaire Tindall, New York 1977; 10.
34. White HS, Wolf HH, Woodhead JH, Kupferberg HJ. The National Institute of health anticonvulsant drugs development programme. Screening for efficacy In: French J, Leppick IE, Dichtes MA, (Eds). Antiepileptic Drug Development: Advances in Neurology, Vol.76, Lippincott-Raven Publishers, Philadelphia 1998; 29-39.
35. Dina M, Lotfy M, Marwa MS, Seham HMH, Sanaa AK. Modulation of PTZ-induced convulsions in rats using topiramate alone or combined with low dose gamma irradiation: involving AKT/m-TOR pathway. Toxicol Mech Met 2021.<https://doi.org/10.1080/15376516.2021.1956032>
36. Dutta A, Handique C, Lahkar M. Evaluation of Anticonvulsant potential of *Bryophyllum pinnatum* in experimental animals. Asian J Biomed Pharmaceut Sci 2015; 5(41); 18-22.<https://doi.org/10.15272/ajbps.v5i41.662>.
37. Sayin U, Cengiz S, Altug T. Vigabatrin as an anticonvulsant against pentyletetrazole seizures. Pharmacol Res 1993; 28: 325-31.<https://doi.org/10.1006/phrs.1993.1134>
38. Rajendra S, JW. Lynch JW, Schofield, PR. The glycine receptor. Pharmacol Ther 1997; 73: 121-146.[https://doi.org/10.1016/S0163-7258\(96\)00163-5](https://doi.org/10.1016/S0163-7258(96)00163-5).
39. Ogbonnia S, VanStaden J, Jager AK, Coker HA. Anticonvulsant effect of *Glyphaea brevis* (Spreng) Moraches leaf extract in mice and preliminary

phytochemical tests. Nig Q J Hosp Med 2003; 13(3-4):62-64.<https://doi.org/10.4314/nqjhm.v13i3-4.12662>.