

A Systematic Review on Molluscicidal Activity of Plants Worldwide Against Intermediate Snail Host of Schistosomes

Kinza Qadeer¹ , Naseer Ali Shah ^{*2} , Haroon Ahmed¹ , Muhammad Kamran³ 

¹COMSATS University Islamabad, Pakistan

²COMSATS University Islamabad, Pakistan

³COMSATS University Islamabad, Pakistan

Article Info	A B S T R A C T
Article type: Review Article Article History: Received: 4 Feb 2025 Revised: 9 May 2025 Accepted: 31 May 2025 Published Online: ✉ Correspondence to: Naseer Ali Shah Email: drnaseer@comsats.edu.pk	Objective: Schistosomiasis affects more than 230 million people worldwide each year. Aquatic snails, which serve as the intermediate hosts, play a key role in transmitting the disease. The present systematic review compiles data on the molluscicidal activity of plants and their phytochemicals to help minimize the incidence of schistosomiasis. Methodology: Online databases such as Google Scholar, PubMed, and ScienceDirect were searched for articles published between 2000 and 2022. In total, 290 articles were identified, of which 193 were included; studies involving snail species not associated with schistosomiasis were excluded. Data such as snail species, plant species and family name, plant parts used, assessment type, and phytoconstituents were analyzed. Results: This review identified 279 plant species with molluscicidal activity, most belonging to the families Euphorbiaceae (9.31%), Solanaceae (7.16%), Asteraceae (6.09%), Leguminosae (5.73%), and Lamiaceae (4.3%). The most frequently used plant parts were leaves (48.74%), followed by seeds (12.9%), fruits (15.05%), roots (14.69%), stems (12.54%), and bark (10%). Conclusion: Moreover, the review highlights the major plant phytochemicals and mechanisms of molluscicidal activity. Plants and their phytochemicals are promising resources for controlling snail species associated with schistosomiasis. Future studies should focus on identifying and isolating active phytochemicals and clarifying their mechanisms of molluscicidal action. Keywords: Schistosomiasis Control, Phytochemicals, Molluscicidal Plants, Biomphalaria, Oncomelania, Intermediate Host
➤ How to cite this paper Qadeer K, Ali Shah N, Ahmed H, Kamran M. A Systematic Review on Molluscicidal Activity of Plants Worldwide Against Intermediate Snail Host of Schistosomes. Plant Biotechnology Persa. 2025; Proof.	

Introduction

Schistosomiasis, commonly known as snail fever, is caused by trematodes belonging to the genus *Schistosoma* [1]. The main *Schistosoma* species that cause schistosomiasis include *S. japonicum*, *S. mansoni*, *S. haematobium*, *S. mekongi*, and *S. intercalatum* [2]. Schistosomiasis is a parasitic disease caused by *Schistosoma* worms, with clinical manifestations varying depending on the species, immune response, and infection duration. Symptoms include fever, fatigue, eosinophilia, abdominal pain, rashes, cough, respiratory issues, muscle and joint pain, nausea, vomiting, and gastrointestinal disturbances. Chronic symptoms include hepatosplenomegaly, intestinal complications, urinary symptoms, neurological symptoms,

immunoreconstitution inflammatory syndrome, and cognitive impairments. Understanding these symptoms can aid in early detection and management, reducing the burden of schistosomiasis on affected individuals and communities [3].

Globally, over 230 million people are infected, with Africa accounting for 95% of cases, highlighting significant regional disparities in disease burden [4]. The regional breakdown of the numbers shows that Africa has the highest prevalence due to a combination of environmental and socioeconomic factors that favour parasite survival. In Uganda, [5] even after treatment, individuals in high-

endemic areas continue to become infected. Schistosomiasis in Asia varies by region, but rural populations in China show a high prevalence [6].

The life cycle of schistosomes is completed in two hosts. Snails, mainly freshwater snails, serve as an intermediate host in which schistosomes undergo asexual reproduction, including the development of miracidia and formation of cercariae (a tailed structure), which upon contact with human skin enter the tissue and develop into adults [1]. The snail species acting as an intermediate host of schistosomes [7] mainly belong to the genera *Biomphalaria*, *Bulinus* and *Oncomelania* (Gastropoda: Pulmonata: Planorbidae) [8,9].

Biomphalaria glabrata, host of *S. mansoni*, is widely found in Africa and South America and sustains transmission in both urban and rural settings. In East and Southeast Asia, *Oncomelania hupensis* is closely linked with *S. japonicum*, particularly in rice paddies where irrigation practices increase exposure risks. *Bulinus truncatus*, the main host of *S. haematobium*, thrives in freshwater habitats across the Middle East and North Africa, playing a central role in urinary schistosomiasis. In contrast, *Austropeplea* species have a more restricted distribution in Sub-Saharan Africa and mainly facilitate livestock-associated *S. bovis* infections. Together, these host-parasite associations underline the ecological diversity of transmission and can help novice readers appreciate regional variations in schistosomiasis epidemiology [10].

In different regions of the world, various control strategies are used to prevent the transmission and outbreak of schistosomiasis [11] and eliminate the disease by disrupting the life cycle of snail species (intermediate host) [12]. Egypt was the first country to control schistosomiasis by using chemical agents against the schistosome-transmitting snail species [13]. The chemicals commonly used as molluscicides include pyrethroids, carbamates, metaldehyde, organophosphates, and niclosamide [14]. Metaldehyde is a widely used, stable molluscicide belonging to the aldehyde group; its application causes snails to secrete an excessive amount of mucus from the body, leading to dehydration and death [15]. Niclosamide is another preferred synthetic molluscicide that restricts the transmission of schistosomes by killing snails or making them inactive [16]. However, these synthetic molluscicides cause environmental pollution.

Recently, the focus has shifted from synthetic molluscicides to plant-based molluscicides because they are economical, easily degradable in the environment and less likely to produce resistance in snails [17]. *Phytolacca dedecandra* (family Phytolaccaceae) is the first plant reported to exhibit molluscicidal activity [18], but its use is now restricted owing to its toxicity to non-target species [11].

An ideal molluscicidal plant must have certain key characteristics. For example, the plant extract should be toxic at extremely low concentrations and non-toxic to non-target species and it needs to remain viable after a period of storage. It is also preferred that the plant species is endemic. Another characteristic of an ideal plant-based molluscicide is that it has a stable structure against environmental fluctuations [19].

This review aims to present worldwide data on plants and their phytochemicals that exhibit molluscicidal activity against intermediate host snail species of schistosomes.

Materials and Methods

Studies regarding the molluscicidal activity of plants against the intermediate host of schistosomes were identified and screened on the basis of certain inclusion and exclusion criteria described later here (Fig 2). Articles were searched using online databases such as Google Scholar, PubMed, and ScienceDirect using keywords such as “plant molluscicide,” “*phytolacca*, natural products against schistosome snail hosts.phytochemicals,” “molluscicidal activity of plants,” “snail sp. causing schistosomiasis” and “plant extracts against snail species.” Only articles available as full texts (n = 290) were downloaded and further screened; of these, duplicates (n = 6), irrelevant studies (n = 4), and studies that did not mention snail species associated with the transmission of schistosomiasis (n = 83) were excluded. The remaining 193 articles were included in this review. Subsequently, data were extracted from 173 articles (Table 1), including information on plant species and family, plant parts used, screened phytochemicals, and their efficacy against snail species.




Figure 1: Study design for the review process.

Results

A total of 193 articles were found eligible to be included in the review, and data from 173 articles related to plant molluscicidal activity against schistosome-transmitting snails were extracted (Table 1).

Table 1: List of plants exhibiting molluscicidal activity against snail species that acts as intermediate hosts of schistosomes (data extracted from articles published between 2000 and 2022)

Snail species	Country	Plant species	Plant Family	Part used	Extraction method	Phytoco nstituent	Assessm ent type	Lethal concentrati on	Findings	Ref.
Biomophalaria alexandrina	Egypt	Moringa oleifera	Moringaceae	Seed	Cerium oxide nano-composite	-----	In-vitro	LC50= 314.5 mg/L, LC90= 386.5 mg/L	Reduced reproductive rate, significant disappearance of cilia and structural deformation in tubular glands	[20]
	Egypt	Colchicum ritchii	Colchicaceae	Flower	Methanolic extract	Tannins, flavonoi ds, saponins , terpenoi ds, steroids, anthraqu inones and alkaloids	In-vitro, In-vivo	LC50 = 35.1 mg/L	Significant increase in micronuclei and 55 % miracidia mortality rate	[21]
	Egypt	Ziziphus spina-christi	Rhamnaceae	Seed	Methanolic extract	Tannins, saponins , alkaloids , flavonoi	In-vitro	LC50 = 108 mg/L	Reduced catalase activity, Increased total antioxidant activity and caused mortality in Daphnia magna	[22]

						ds, phenols and glycosid es				
	Egypt	Carica papaya	Caricaceae	Seed	Methanolic extract	Tannin and saponin	In-vitro	LC50= 138.5 mg/L	Increased catalase activity, Slight increase in total antioxidant activity and caused mortality in Daphnia magna	
	Egypt	Origanum majorana	Lamiaceae	Leaves	Aqueous extract	Zingiber ene, nerolidol , cubenol, eudesme nol, cedren, butanon e, capsaicin , gingerol, stigmast erol and nonivam ide	In-vitro	LC50= 42 mg/ml	Swelling of a cephalopodal mass and hemorrhagic blisters on the foot sole	[23]
	Egypt	Ziziphus spina-christi	Rhamnaceae	Leaves	Aqueous extract	Cyclooct asiloxan e, hydrochi non, morphin	In-vitro	LC50= 43 mg/ml	Reduced miracidal activity and 100 % cercarial activity at 6.25 mg/ml	[23] [22] [22] [22] [22]

						an, tricyclou ndecan, spathule nol, imidazol e, megastig matrieno ne, guanosin e, desulpho sinigrin, phenant hreneme thanol and thienopy ridine				[22] [22]
	Egypt	Salvia fruticosa	Lamiaceae	Leaves	Aqueous extract	Quinolin ol, tetrasilo xane, androsta dienol, purinol, spiroste n,	In-vitro	LC50= 69 mg/ml	Less effective against miracidial activity and 100 % cercarial activity at 100 mg/ml	

						tetradecanoic acid, benzene methanol and cystathionine and terphenyl				
	Egypt	Ziziphus spina christi	Rhamnaceae	Leaves	Ethanolic extract	-----	In-vitro	LC50= 108.7 mg/l	Increased apoptosis and condensed zygote	[24]
	Egypt	Moringa oleifera	Moringaceae	Leaves	Ethanolic extract	-----	In-vitro	LC50= 209.4 mg/l	Increased apoptosis and condensed zygote	
	Egypt	Tecoma stans	Bignoniaceae	Leaves	Ethanolic extract	-----	In-vitro	LC50= 256 mg/l	Increased apoptosis and condensed zygote	
	Egypt	Carica papaya	Caricaceae	Leaves	Methanol, ethanol and butanol extract	----- -	In-vitro	LC50 (mg/L) against cercariae For methanol= 2; For ethanol=	65-86 % mortality rate	[25]

								20 and for butanol= 4		
	Egypt	Anagallis arvensis	Primulaceae	Whole plant	Ethanollic extract	----- --	In-vitro	LC50= 1.76 ppm ; LC90= 3.16 ppm	Reduction in protein content, denatured hermaphrodite gland, decreased amount of connective tissues and 100 % mortality rate in case of LC90 after 4 hr	[26]
	Egypt	Anagallis arvensis	Agavaceae	Whole plant	Methanolic extract	----- ---	In-vitro	LC50= 45 ppm LC90= 60 ppm	Extensive vacuolation in cells of digestive system and 100 % mortality at 70 ppm	[27]
	Egypt	Viburnum tinus	Caprifoliaceae	Whole plant	Methanolic extract	----- ---	In-vitro	LC50= 38 ppm LC90= 59 ppm	90 % mortality at 60 ppm, deformed gonadal cells, reduced number of sperms and degenerated cercariae	
	Egypt	Anagallis arvensis	Primulaceae	Leaves	Aqueous extract	----- -	In-vitro	LC50= 37.9 mg/l LC90= 48.3 mg/l	Decreased reproductive rate, deformation in hermaphrodite gland cells and loss of connective tissues in gonadal cells	[28]
	Egypt	Albizia anthelmintica	Leguminosae	Leaves	-----	Saponin	In-vitro	LC50 = 17.6 ppm LC90 = 25.3 ppm	Decrease in granulocytes and hyalinocytes while increased amoebocytes	[29]

	Egypt	Callistemon citrinus	Myrtaceae	Leaves	Methanol extract	----- -	In-vitro	LC90= 37.8 ppm	Loss of hyaline substance, loss of connective tissues and formation of vacuoles	[30]
	Egypt	Callistemon citrinus	Myrtaceae	Leaves	Methanolic extract	----- -	In-vitro	LC50= 22.3 ppm LC90 = 37.8 ppm	54.2 % reduction in reproductive rate, crumpling of young oocytes and 56% reduction of testosterone	[31]
	Egypt	Zingiber officinale	Zingiberaceae	Root	Methanolic extract	----- -	In-vitro	LC50= 43.8 ppm LC90 = 74.9 ppm	60 % reduction in reproductive rate, atrophy of nuclei and degeneration of necrotic cells	
	Egypt	Asparagus densiflorus	Asparagaceae	Leaves	Aqueous extract	----- -	In-vivo	LC50 = 75 ppm LC90 = 102 ppm	Caused atrophy in gonadal cells, reduced reproductive rate and 100% mortality	[32]
	Egypt	Oreopanax guatemalensis	Araliaceae	Leaves	Aqueous extract	----- -	In-vivo	LC50 = 160 ppm LC90 = 188 ppm	100 % mortality rate at 1000 mg/L within 24 h	
	Egypt	Juniperus horizontalis	Cupressaceae	Aerial parts	Methanolic extract	Thujone,	In-vitro	LC50 = 38.9 ppm	Proved toxic against snail sp.	[33]

						D-limonene, β-myrcene, sabinene and methyl citronellate				
	Egypt	Juniperus communis	Cupressaceae	Aerial parts	Methanolic extract	Sabinene, β-myrcene, D-limonene, thujone and terpinen-4-ol	In-vitro	LC50 = 22.9 ppm	Proved toxic against snail sp.	
	Egypt	Cestrum purpureum	Solanaceae	Whole plant	Methanolic extract	ursolic acid and 3-O-β-D-glucopyranosyl-(	In-vivo	LC50= 72 ppm LC90 = 120 ppm	90 % mortality rate after 24h	[34]

						1→4)-α-L-rhamnopyranosyl-2 and 19-dihydroxy-epihederagenin-28-O-1-[26-(3-acetyl-4-methoxycinnamoyloxyhexacosanoyl)] glycerol				
	Egypt	Agave angustifolia	Agavaceae	Whole plant	Aqueous suspension	----- -	In-vivo, In-vitro	LC90= 120 ppm	Decreased egg laying capacity and survival rate	[35]
	Egypt	Pittosporum tobira	Pittosporaceae	Whole plant	Aqueous suspension	----- -	In-vivo, In-vitro	LC90= 120 ppm	Decrease in cercaricidal mortality by 81% and miracidia by 87 %	
	Egypt	Euphorbia peplodes	Euphorbiaceae	Aerial parts	Methanolic extract	Quercetin, luteolin, gallic acid, β-amyrin,	In-vivo	LC90= 30 ppm	Strong molluscicidal activity	[36]

						P-hydroxy benzoic acid, quercetin-3-O- β -D-glucopyranoside and kampferol-3-O- β -D-glucopyranoside				
	Egypt	Callistemon viminalis	Myrtaceae	Leaves, fruit and bark	Methanolic extract	----- -	In-vivo	LC50 (ppm): Fruit=6.2, Bark=32, Leaves =40	63.3 % mortality rate due to leaves extract, 66.7% due to bark and 100% due to leaves	[37]
	Egypt	Eucalyptus globulus, Melaleuca styphelioides	Myrtaceae	Leaves	Methanolic extract	Eucalbanin B, tellimagrandin I, pentagalloyl glucose, casuarinin and 1,2,3,4,6-Penta-O-galloyl-b-D-	In-vitro	LC50= 165.6 ppm LC90= 245.5 ppm	25 % reduction in protein content, 100 % cercaricidal mortality, 21 % increase in acid phosphatase and 113 % increase in aspartate aminotransferase	[38]

						glucopyranose				
	Egypt	Citrus limon	Rutaceae	Peel	Aqueous extract	----- --	In-vitro	LC50 = 744.17 ppm LC90 = 1821.25 ppm	58 % reduction in hatchability rate, necrosis in digestive glands and significant increase in protein content	[39]
	Egypt	Origanum syriacum	Lamiaceae	Seed	Ethanol extract	----- --	In-vitro	B. alexandrina LC50 = 226.8 ppm LC90 = 422.9 ppm	Chromosomal aberrations	[40]
	Egypt	Yucca desmettiana	Asparagaceae	Leaves	Methanolic extract	Saponin	In-vivo	LC50 = 68 mg/L LC90 = 96 mg/L	Significant molluscicidal activity	[41]
	Egypt	Haptophyllum tuberculatum	Rutaceae	Aerial parts	Chloroform and methanolic extracts	Steroids	In-vitro	LC50 = 16 ppm	100 % mortality rate at 25 ppm with excessive mucous production, increased hemolysis and decreased progesterone level	[42]

	Egypt	<i>Datura stramonium</i>	Solanaceae	Leaves	Dry powder	----- -	In-vitro	LC50= 25.8 ppm LC90= 40.4 ppm	60 % mortality rate, increased alanine transaminase and alkaline phosphatase in haemolymph	[43]
	Egypt	<i>Sesbania sesban</i>	Leguminosae	Leaves	Dry powder	----- -	In-vitro	LC50= 51.0 ppm LC90= 62.4 ppm	80 % mortality rate and decreased total protein content	
	Egypt	<i>Solanum siniacum</i>	Solanaceae	Leaves	Ethanolic, aqueous, methanolic and chloroform extract	----- -	In-vitro	LC50 = 14.8 ppm	88 % reduction in reproductive rate, 100 % increase in glucose and 80 % in lactate and significant decrease in glycogen, pyruvate and total protein amount	[44]
	Egypt	<i>Artemisia judaica</i> L	Asteraceae	Leaves	Ethanolic, aqueous, methanolic and chloroform extract	----- -	In-vitro	LC50 = 38 ppm	Increase in protease quantity but decreased lactic dehydrogenase and cytochrome oxidase	
	Egypt	<i>Dalbergia sissoo</i>	Leguminosae	Fruits, leaves, root and stem bark	Ethanolic extracts	----- -	In-vivo	LC50= 10.8 ppm LC90= 20.4 ppm	Mainly toxic at pre-hatch stage and 100 % mortality rate	[45]

	Egypt	<i>Furcraea selloa marginata</i>	Agavaceae	Leaves	Aqueous suspension	----- -	In-vitro	LC50 =53.66 ppm LC90 =84.35 ppm	30 % mortality rate after 24 hr and 100 % miracidial mortality rate	[46]
	Egypt	<i>Euphorbia milii</i>	Euphorbiaceae	Latex	Aqueous solution	----- -	In-vitro	LC50= 19 ppm LC90= 38 ppm	Proved toxic against snail sp.	[47]
	Egypt	<i>Meryta denhamii</i>	Araliaceae	Flower	Methanolic extract	Oleonolic acid-3-O- α -arabinopyranoside, oleonoic acid-3-O- β -glucopyranoside and hedragenin-3-O- α -rhamnopyranoside	In-vivo	LC50=85 mg/L	Increased glucose level and decreased glycogen and protein level in tissues	[48]

	Egypt	Guayacum officinalis	Zygophyllaceae	Whole plant	Methanolic, aqueous and ethanolic extract	----- -	In-vitro	LC50= 120 ppm LC90= 210 ppm	Decreased acetylcholinesterase and Succinate dehydrogenase	[49]
	Egypt	Atriplex stylosa	Chenopodiaceae	Whole plant	Methanolic, aqueous and ethanolic extract	----- -	In-vitro	LC50= 94 ppm LC90= 180 ppm	36 % decreased glycogen content and 53 % increased glucose level	
	Egypt	Euphorbia splendens	Euphorbiaceae	Whole plant	Methanolic, aqueous and ethanolic extract	----- -	In-vitro	LC50= 40 ppm LC90= 73 ppm	Effectively decreased the number of juveniles and adult	
	Egypt	Citrus reticulata	Rutaceae	Fruit	Peel extract	----- -	In-vivo	LC50= 163.79 ppm	95 % mortality rate	[50]
	Egypt	Euphorbia paralias	Euphorbiaceae	Aerial parts	Methanolic extract	Diterpenes (Paralianses, segetanes and jatrophe-nes)	In-vivo	-----	100 % mortality rate	[51]
	Egypt	Commiphora molmol	Bruseraceae	Stem (Gum resin)	Aqueous solution	Terpenes and eugenol	In-vivo	-----	67 % mortality rate	[52]

	Egypt	Commiphora molmol	Bruseraceae	Whole plant	Oil extract	----- -	In-vivo	LD50= 155 ppm LD90= 195 ppm	100 % mortality rate	[53]
	Egypt	Yucca elephantipes	Agavaceae	Leaves	Methanolic extract	saponin	In-vivo	LC90= 4 ppm	Strong molluscicidal activity	[54]
	Egypt	Ambrosia Maritima	Asteraceae	Whole plant	Dry powder	----- ---	In-vivo	-----	95 % decrease number of snails in first month	[55]
	Egypt	Azolla pinnata	Salviniaceae	Whole plant	Dry powder	----- ---	In-vivo	-----	94 % decrease number of snails in first month	
	Egypt	Commiphora molmol	Burseraceae	Whole plant	Oil and resin extract	----- ---	In-vivo	LC50= 4 ppm	100 % mortality rate of both snail sp.	[56]
	Saudi Arabia	Rosmarinus officinalis	Lamiaceae	Whole plant	Pet-ether and ethanolic extracts	----- -	In-vivo	LC50= 236.81 ppm	85 % mortality rate	[57]
	Italy	Vitex trifolia	Verbenaceae	Leaves, stem and seed	chloroform, n-hexane and ethanolic extract	Platanic acid , β -sitosterol, ursolic acid acetate and ursolic acid acetate	In-vivo	LC50= 26.42 mg/l	92 % mortality rate at 32.01 mg/l	[58]

Biomphalaria pfeifferi	Zimbabwe	Cucurbita maxima	Cucurbitaceae	Seed	Aqueous and crude ethanolic extracts	----- -	In-vivo	LC50 (mg/ml) for: Aqueous extract = 0.002, Ethanolic extract = 0.002	100 % mortality rate at 2 mg/mL in both snails	[59]
	Nigeria	Balanites aegyptiaca	Balanitaceae	Leaves and fruit	Aqueous extract	Alkaloids , saponins , flavonoids, tannins, phenols and steroids	In-vitro	-----	90 % mortality rate at 50 µg/ml	[60]
	Sudan	Combretum glutinosum Perr. ex DC	Combretaceae	leaves	Ethanolic extract	Cumarins, tannins, flavonoids, saponins and	In-vivo	LC50 at 24 hr: 117.57 LC90 at 24 hr: 220.84	Possess strong molluscicidal ability	[61]

						triterpenes				
	Sudan	Hyphaene thebaica (L.) Mart	Arecaceae	Male Inflorescences	Ethanolic extract	Cumarins, tannins and saponins	In-vivo	LC50 at 24 hr: 158 LC90 at 24 hr: 228	Caused mortality at low concentration	
	Sudan	Indigofera oblongifolia Forssk	Leguminosae	Aerial parts	Ethanolic extract	Cumarins, tannins, saponins, triterpenes and steroids	In-vivo	LC50 at 24 hr: 158.11 LC90 at 24 hr: 228.11	Mortality rate was more than 90 %	
	Sudan	Rhynchosia minima (L.) DC	Leguminosae	Aerial parts	Ethanolic extract	Cumarins, tannins, flavonoids and saponins	In-vivo	LC50 at 24 hr: 158.11 LC90 at 24 hr: 228.11	Caused mortality at low concentration	

	Sudan	<i>Solanum dubium</i> L.	Solanaceae	Fruit	Ethanol extract	Cumarins, tannins, flavonoids, saponins, alkaloids and steroids	In-vivo	LC50 at 24 hr: 153.02 LC90 at 24 hr: 226.62	90 % mortality at low concentration	
	Sudan	<i>Pulicaria crispa</i>	Asteraceae	Leaves, stem and root	Powder	----- -	In-vivo	LC50(ppm): Leaves=616.6 Stem=616.6 Root=645.65 LC90(ppm): Leaves=776.25 Stem=812.83 Root=812.83	Sluggish and disoriented behavior	[62]

	Ethiopia	<i>Achyranthes aspera</i>	Amaranthaceae	Leaves	Hydro-ethanolic extract	Flavonoids, saponins, alkaloids and terpenoids	In-vivo	In immersion: LC50(ppm) = 20.37 LC90 (ppm) = 46.84; In immersion: LC50 = 3.10 ppm LC90 = 11.08 ppm	30-40 % mortality rate after 24 hr at 100 ppm	[63]
	Sudan	<i>Solenostemma argel</i>	Asclepiadaceae	Leaves	Aqueous extract	----- -	In-vivo	For egg masses, LD50 = 2921.3 ppm LD95 = 3515 ppm; For neonates, LD50 = 368 ppm, LD95 = 1699 ppm	50 % mortality rate of eggs and 95 % of neonates	[64]

	Kenya	<i>Aloe secundiflora</i>	Asphodelaceae	Leaves	Aqueous extract	----- -	In-vivo	-----	40 % mortality rate after 24 h	[65]
	Kenya	<i>Balanites aegyptiaca</i>	Zygophyllaceae	Root	Aqueous extract	----- -	In-vivo	-----	36 % mortality rate after 24 h	
	Kenya	<i>Aspilia pluriseta</i>	Asteraceae	Leaves	Aqueous extract	----- -	In-vivo	-----	50 % mortality rate after 24 h	
	Kenya	<i>Psidium guajava</i>	Myrtaceae	Root and leaves	Aqueous extract	----- -	In-vivo	-----	20 % mortality rate after 24 h	
	Kenya	<i>Azadirachta indica</i>	Meliaceae	Root and leaves	Aqueous extract	----- -	In-vivo	-----	58 % mortality rate after 24 h	
	Sudan	<i>Solenostemma argel</i>	Apocynaceae	Leaves	Aqueous extract	Alkaloids , terpenoids, flavonoids, saponins and tannins	In-vivo	LD50= 0.103 ppm LD95= 0.187 ppm	Behavioral changes	[66]
	Nigeria	<i>Balanites aegyptiaca</i>	Zygophyllaceae	Stem and bark	Aqueous solution	2-methoxy-4-vinylphenol, benzaldehyde, phenol, 2-	In-vivo	LD50 (mg/L): For <i>B. pfeifferi</i> juvenile=1.4	100 % mortality rate of both snail species at 40mg/L	
										[67]

						methoxy -4-(2- propenyl)- acetate, ethanon e, 1- (4hydro xphenyl) -, alpha- d-6,3- furanose, 4-(3- hydroxy- 1- propenyl)-2- methoxy phenol, hexadec anoic acid, 2- butanon e, 9- octa decanoic acid, octadeca noic acid, 13- octadece nal, tetradec anoic, 9- octadece namide and di-n-		For B. pfeifferri adult=1.6 LD90 (mg/L): For B. pfeifferri juvenile=4. 8 For B. pfeifferri adult=4.2		
--	--	--	--	--	--	--	--	---	--	--

						octylphalate				
	Mali	<i>Glinus oppositifolius</i>	Aizoaceae	Whole plant	Crude extract	12-Glucosyl-GO1	In-vivo	LC50 of Butanol extract= 91.7 ppm; LC50 of Ethanolic extract= 116.6 ppm	100 % mortality rate of both snail species at 200ppm	[68]
	Ethopia	<i>Glinus lotoides</i>	Molluginaceae	Fruit	Aqueous and crude extract	----- -	In-vitro	LC50= 47.1 mg/L LC90= 56.96 mg/L	20 % mortality rate of cercariae, and 100 % snail mortality at 37.5-80 mg/L	[69]
	Ethopia	<i>Balanites aegyptiaca</i>	Zygophyllaceae	Seeds, mesocarp, fruit and endocarp	Aqueous extract	----- -	In-vivo	LC90 (mg/L): Seed= 77.70, Endocarp= 120.04, Mesocarp= 89.50, Fruit=99.55	90 % mortality rate	[70]

	Kenya	Entada leptostachya	Leguminosae	Root	Ethyl acetate, aqueous and methanol extract	Tannins, glycosides, Saponins, flavonoids and triterpenes	In-vitro	LD50 (mg/l): Juvenile= 30.12 Adult= 40.93	100 % mortality rate on juveniles and 95 % on adult	[71]
	Kenya	Azadirachta indica	Meliaceae	Leaves	Ethyl acetate, aqueous, methanol extract	Alkaloids, glycosides, saponins, flavonoids and triterpenes	In-vitro	-----	Only 5 % mortality	
	Nigeria	Chromolaena odorata	Asteraceae	Leaves	Aqueous and ethanolic extract	----- --	In-vivo	LC50 (ppm) for Aqueous extract: Eggs= 65.75 Juveniles= 75.59 Adults= 217.57 LC50 (ppm) for	100 % mortality rate after 24 h	[72]

								Ethanollic extract: Eggs= 44.03 Juveniles= 44.68 Adults= 88.04		
	Nigeria	Ficus exasperata	Moraceae	Root, bark, seed and leaves	Aqueous and ethanolic extracts	Flavonoids, alkaloids, tannins, saponins, anthraquinones and cardenolides	In-vitro	LC50= 0.3187 ppm LC90= 0.7555 ppm	Mitochondrial eruption, degeneration of epithelial cells and 100 % mortality rate	[73]
	Nigeria	Cymbopogon citratus	Poaceae	Leaves	Ethanol and aqueous extract	----- -	In-vivo	LC50 for aqueous extract= 140.74 ppm LC50 for ethanolic extract= 254.92 ppm	100 % mortality in ethanol extract	[74]

	Nigeria	Chromolaena odorata	Asteraceae	Leaves	Ethanollic and aqueous extract	----- -	In-vivo	Aqueous extract: LC50 =65.75 ppm LC90=139.54 ppm Ethanollic extract LC50 =44.03 ppm LC90 =119.3 ppm	100 % mortality rate at 100 ppm	[75]
	Sudan	Acacia seyal	Mimosaceae	Bark	Methanollic extract	----- -	In-vitro	IC50 (ppm): At 24 h =80.79, At 48 h =34.33	100 % mortality at 62.5 ppm after 48 h	[76]
	Nigeria	Piper guineense	Piperaceae	Fruit	Aqueous and ethanollic extract	----- -	In-vivo	For ethanollic extract: LC50 = 0.10 mg/L LC90= 0.9 mg/L	90% mortality rate at 9 mg/L	[77]

								For aqueous extract: LC50 = 5.0 mg/L LC90= 8.5 mg/L		
	Nigeria	Anonna senegalensis	Annonaceae	Leaves and root	Ethanolic extract	----- -	In-vivo	For leaves: LC50 = 88.93 mg/l LC90= 285.44 mg/l For roots: LC50 = 861.38 mg/l LC90= 2 379.80 mg/l	100 % mortality rate of adult snail at 500 mg/l and 75% death rate of egg masses at 250 mg/l	[78]
	Nigeria	Anogeissus leiocarpus	Combretaceae	Leaves and root	Ethanolic extract	----- -	In-vivo	For leaves: LC50 = 1 213.39 mg/l LC90= 3 263.00 mg/l	65 % mortality rate of adult snail at 1000 mg/l and 90% death rate of egg masses at 4000 mg/l	

								For roots: LC50 = 653.60 mg/l LC90= 1 503.03 mg/l		
	Nigeria	Crotalaria retusa	Leguminosae	Leaves and root	Ethanolic extract	----- -	In-vivo	For leaves: LC50 = 73.84 mg/l LC90= 176.94 mg/l For roots: LC50 = 290.35 mg/l LC90= 714.89 mg/l	70 % mortality rate of adult snail at 500 mg/l, 69 % death rate of egg masses at 1000 mg/l	
	Nigeria	Dalbergia sissoo	Leguminosae	Fruit, stem and bark	Ethanolic extract	----- -	In-vivo	For fruits: LC50 = 30.74 mg/l LC90= 74.33 mg/l For bark:	85 % mortality rate of adult snail at 500 mg/l and 81 % death rate of egg masses at 500 mg/l	

								LC50 = 399.86 mg/l LC90= 114.84 mg/l		
	Nigeria	Vernonia amygdalina	Asteraceae	Leaves and root	Ethanolic extract	----- -	In-vivo	For leaves: LC50 = 1913.45mg/l LC90= 6241.61 mg/l For roots: LC50 = 917.79 mg/l LC90= 2384.78 mg/l	85 % mortality rate of adult snail at 2000 mg/l and 89 % death rate of egg masses at 4000 mg/l	
	Nigeria	Carica papaya	Caricaceae	Leaves	Ethanolic extract	----- -	In-vivo	LC50 B. pfeifferi=2716.3 ppm LC90= 4515.9 ppm	50 % mortality rate	[79]

	Nigeria	<i>Terminalia catappa</i>	Combretaceae	Leaves	Ethanollic extract	----- -	In-vivo	LC50 = 864.1 ppm LC90= 1222.8 ppm	100 % mortality rate	
	Sudan	<i>Cymbopogon nervatus</i>	Poaceae	Leaves	Essential oils	----- -	In-vivo	LD50 = 213.099 ppm	100 % mortality rate at 400 ppm	[80]
	Sudan	<i>Boswellia papyrifera</i>	Burseraceae	Leaves	Essential oils	----- -	In-vivo	LD50 213.31 ppm	90 % mortality rate at 350 ppm	
	Nigeria	<i>Dalbergia sissoo</i>	Leguminosae	Fruit, root, stem and leaves	Ethanollic and aqueous extracts	----- -	In-vitro	LC90 =89.29 mg/l	93 % death rate of snail eggs at 500 mg/l	[81]
	Nigeria	<i>Dalbergia sissoo</i>	Leguminosae	Fruit, root, leaves and stem	Aqueous and ethanollic extract	----- -	In-vitro	-----	Un-hatched embryo death	[82]
	South Africa	<i>Barringtonia racemosa</i>	Lecythidaceae	Fruit and seed	Aqueous extract	----- -	In-vitro	LD50 for fruit= 521.48 ppm LD50 for seed= 556.23 ppm	More toxic effects are produced by fruit extract	[83]

	Sudan	<i>Azadirachta indica</i>	Meliaceae	Leaves and seeds	Aqueous extract	----- -	In-vivo	-----	75 % decrease in snail species after 5 days	[84]
	Mali	<i>Cussonia barteri</i>	Araliaceae	Roots	Methanolic and aqueous extract	----- -	In-vitro	-----	Extract proved toxic against snail sp.	[85]
	Mali	<i>Glinus oppositifolius</i>	Aizoaceae	Whole plant	Methanolic and aqueous extract	----- -	In-vitro	-----	Extract proved toxic against snail sp.	
	Mali	<i>Lannea velutina</i>	Anacardiaceae	Leaves	Methanolic and aqueous extract	----- -	In-vitro	-----	Extract proved toxic against snail sp.	
	Kenya	<i>Solanum aculeastrum</i>	Solanaceae	Root, bark and fruit	Methanolic extract	----- -	In-vivo	-----	100 % mortality rate	[86]
	Saudi Arabia	<i>Jatropha glauca</i>	Euphorbiaceae	Leaves	Chloroform extract	----- -	In-vivo	LD50= 66.7 ppm LD90= 114 ppm	Decreased efficacy with increased time	[87]
	Saudi Arabia	<i>Euphorbia helioscopia</i>	Euphorbiaceae	Leaves	Methanolic extract	----- -	In-vivo	LD50= 50.8 ppm LD90= 68.2 ppm	Decreased efficacy with increased time	
	Saudi Arabia	<i>Euphorbia schimperiana</i>	Euphorbiaceae	Stem	Methanolic extract	----- -	In-vivo	LD50= 34 ppm	100 % mortality rate	

								LD90= 41.7 ppm		
Biomphalaria glabrata	Egypt	Origanum syriacum	Lamiaceae	Seed	Ethanolic extract	----- -	In-vitro	LC50= 215.2 ppm LC90= 415.7 ppm	Chromosomal aberration in both snail species	[40]
	Nigeria	Lycopersicon esculentum	Solanaceae	Seed	Methanolic extract	----- -	In-vivo	LC50 = 873.6 ppm	78 % mortality at 1000 µg/ml	[88]
	Nigeria	Blighia unijugata	Sapindaceae	Stem, fruit, seed and leaves	Ethanolic extract	Saponin	In-vivo	LC50= 7.6 µg/ml	100 % mortality rate at 1000 ppm	[89]
	Angola	Euphorbia conspicua	Euphorbiaceae	Latex	Methanolic extract	----- -	In-vivo	LC100 = 1 µg/mL	Toxic against the snail sp.	[90]
	Mali	Cussonia barteri	Araliaceae	Roots	Methanolic and aqueous extract	----- -	In-vitro	-----	Extract proved toxic against snail sp.	[85]
	Mali	Glinus oppositifolius	Aizoaceae	Whole plant	Methanolic and aqueous extract	----- -	In-vitro	-----	Extract proved toxic against snail sp.	
	Mali	Lannea velutina	Anacardiaceae	Leaves	Methanolic and aqueous extract	----- -	In-vitro	-----	Extract proved toxic against snail sp.	
	Thailand	Solanum xanthocarpum	Solanaceae	Fruit	Ethanolic crude extract	----- -	In-vivo,	LC50= 163.85 mg/l LC90=	Extract proved toxic against snail	[91]

							In-vitro	219.33 mg/l		
	Pakistan	Euphorbia cauducifolia	Euphorbiaceae	Latex	DMSO suspension	Deoxyphorbol Esters	In-vivo	LC50=12.3 ng/ml	Induces imbalance in nutrient and water absorption	[92]
	Colombo	Croton floribundus	Euphorbiaceae	Leaves and bark	Ethanollic and methanolic extract	Kaurenoic acid	In-vivo	LC50=4.2 µg/mL LC90=11.5 µg/mL	High molluscicidal activity	[93]
	Japan	Ambrosia maritima	Asteraceae	Root	Chloroform extract	Ambrosin and Pentacyclic	In-vivo	LC50 = 0.37ppm	80 % molluscicidal activity after 72h	[94]
	Brazil	Parkia pendula	Leguminosae	Seed	Saline extract	----- -	In-vitro, In-vivo	LC50 (mg/mL) values: Blastula= 237.19 Gastrula= 244.62 Adult= 4.99	Proved toxic at 3.0, 4.0 and 5.0 mg/mL, decrease in defense cells and detection of nuclear damage	[95]

	Brazil	<i>Dysphania ambrosioides</i> (L.)	Amaranthaceae	Aerial parts	Essential oil	α -Terpinene, ascaridole, thymol, carvacrol, isoascaridole, p-cymene, limonene and γ -terpinene	In-vitro, In-vivo	LC50= 25.2 μ g/mL	100 % mortality rate at 50 μ g/mL, non-toxic to <i>Artemia salina</i> larvae and <i>Danio rerio</i> fish	[96]
	Brazil	<i>Persea Americana</i> Mill.	Lauraceae	Stem and bark	Ethanolic extract	Quercetin, flavonoid, anthraquinone heterosides, coumarins and tannins	In vivo, In vitro	LC50 (ppm): For embryo =27.06 For newly hatched= 30.6 For adult =55.55	Increased embryo mortality at 144 h, 100 % mortality in adult, lethargy and shell reduction after 96 h	[97]
	Brazil	<i>Pimenta dioica</i>	Myrtaceae	Leaves	Essential oil	Octenol, mirceno, limonene, linalol, terpinol, chavicol	In-vivo	LC50= 18.62 mg/L	90 % mortality rate after 72 h at 75 mg/L	[98]

						and eugenol				
	Brazil	Citrus limon	Rutaceae	Fruit peel	Essential oil	Limonene, sabinene, β -pinene, γ -terpinene, β -citral, α -citral, α -pinene and β -Bisabolene	In-vivo	-----	100 % mortality at 100 mg/L	[99]
	Brazil	Manilkara subsericea	Sapotaceae	Leaves	Ethanollic crude extract and ethyl acetate fraction	Myrcetin, quercetin and ursolic acid	In-vivo	LD50 (μ g/mL): For Ethanollic crude extract= 118.7 For ethyl acetate	53 % mortality rate at 250 μ g/mL	[100]

								fraction= 23.41		
	Brazil	<i>Euphorbia umbellata</i>	Euphorbiaceae	Latex	Latex aqueous dilutions	Triterpenes	In-vivo	LC50=1.36 mg/L LC90 =3.69 mg/L	90 % mortality rate after 24 h,	[101]
	Brazil	<i>Anadenanthera colubrina</i>	Planorbiidae	Bark	Crude extract	Terpenoids, tannins and phenols	In-vivo	LC50 For adult= 0.155 mg/mL For embryo= 0.692 mg/mL	100 % mortality of adult snail and embryo at 0.5, 1.0 mg/mL respectively	[102]
	Brazil	<i>Moringa oleifera</i> Lam.	Moringaceae	Flower	Aqueous extract	Xanthone, saponin, tannin, flavanol, polyphenol and flavones	In-vivo	LC50= 2.37mg/mL	95 % mortality rate at 4 mg/mL	[103]
	Brazil	<i>Jatropha elliptica</i>	Euphorbiaceae	Root	Ethanollic extract	diethyl 4-phenyl-2,6-dimethyl-3,5-	In-vivo	LC90=36.43 µg/mL	100 % mortality rate of schistosomes and snail	[104]

						pyridine dicarboxylate				
	Brazil	Schinopsis brasiliensis	Anacardiaceae	Stem, bark	Hydroethanol extract	Xanthones, flavanones and tannin	In-vivo	LC90=73 µg/ml	60 % morality rate at 50 µg/ml	[105]
	Brazil	Jatropha gossypifolia	Euphorbiaceae	Stem, leaves and fruit	Hydroalcoholic extract	Alkaloids, tannins, triterpenoids, saponins and steroids	In-vivo	-----	100 % mortality rate at 25 ppm, significant reduction in oviposition ability and complete repression of feeding ability	[106]
	Brazil	Moringa oleifera	Moringaceae	Seed	Aqueous extract	----- --	In-vivo	LC50 =0.419 g/l; LC90 =1.021 g/l	Toxic for Biomphalaria glabrata	[107]
	Brazil	Piper tuberculatum	Piperaceae	Stem, leaves, fruit and root	Methanolic extract	Piplartine	In-vitro	LC50 (µg/ml): Root= 20.28 Stem=200 Leaves= 310.27	100 % mortality rate of blastula, gastrula and adult	[108]

								Fruit= 126.27		
	Brazil	<i>Cymbopogon winterianus</i>	Poaceae	Leaves	Essential oil	Geraniol, citronellol, citronellal and elemol	In-vivo	LC50=54.0 mg/L, LC90=97.0 mg/L,	90 % mortality rate after 72 h	[109]
	Brazil	<i>Porophyllum ruderale</i>	Asteraceae	Flower and leaves	Essential oil	β -pineno, mircene, (E)- β -ocimeno and (Z)- β -ocimeno,	In-vivo	LC50 = 774.82ppm LC90 = 812.43ppm	Proved toxic at less than 1000 ppm	[110]
	Brazil	<i>Euphorbia milii</i>	Euphorbiaceae	Latex	Aqueous solution	----- --	In-vivo	LC50= 0.7 μ l	Decreased glycogen content, increased glucose and total protein content	[111]
	Brazil	<i>Euphorbia milii</i>	Euphorbiaceae	Latex	Aqueous solution	----- --	In-vivo	EC50 = 2040 μ g/L	Increased malformation in embryos	[112]
	Brazil	<i>Lippia gracilis</i>	Solanaceae	Leaves	Essential oil	Thymol, methyl-thymol and p-cymene	In-vivo	LC50 = 62.2 ppm LC90= 82.8 ppm	Proved highly lethal to snail sp.	[113]

	Brazil	Turnera ulmifolia	Turneraceae	Leaves	Hydroalcoholic extract	Glycosides, flavonoids, alkaloids, tannins and steroids	In-vivo	-----	No activity against snail sp.	[114]
	Brazil	Solanum asperum	Solanaceae	Fruit	Methanolic extract	Solanandaine and solamargine	In-vivo	LC50= 25.1 µg/mL LC90= 44.1 µg/mL	90 % mortality rate after 24h	[115]
	Brazil	Ocotea bracteosa	Lauraceae	Stem and bark	Essential oil	Cadinene, ledene and globulol	In-vivo	LC90 = 8.3 µg/mL	Proved toxic against the snail sp.	[116]
	Brazil	Euphorbia splendens	Euphorbiaceae	Latex	Aqueous stock solution	----- --	In-vitro	LD50 = 1 mg/l LD90 = 2.3 mg/l	Increased protein conc. and decreased glycogen content	[117]
	Brazil	Solanum stipulaceum	Solanaceae	Stem	Aqueous extract	----- --	In-vivo	LC50= 45.2 µg/mL LC90= 56.0 µg/mL	Toxic against the snail sp.	[118]

	Brazil	<i>Solanum diamantinense</i>	Solanaceae	Aerial parts	Methanolic extract	----- --	In-vivo	LC50= 33.36 µg/mL LC90= 52.8 µg/mL	High molluscicidal activity against snail sp.	[119]
	Brazil	<i>Solanum paludosum</i>	Solanaceae	Fruit	Methanolic extract	----- --	In-vivo	LC50= 48.07 µg/mL LC90= 82.86 µg/mL	High molluscicidal activity against snail sp.	
	Brazil	<i>Stryphnodendron adstringens</i>	Mimosoideae	Leaves and bark	Ethanolic extract	Flavonoids	In-vivo	-----	90 % mortality rate at 50 ppm	[120]
	Brazil	<i>Stryphnodendron polyphyllum</i>	Mimosoideae	Leaves and bark	Ethanolic extract	Flavonoids	In-vivo	-----	70 % mortality rate of adult snail	
	Brazil	<i>Dimorphandra mollis</i>	Mimosoideae	Leaves and bark	Ethanolic extract	Flavonoids, coumarins and tannins	In-vivo	-----	10 % mortality rate at 100 ppm after 48 h	
	Brazil	<i>Caryocar brasiliensis</i>	Caryocaraceae	Leaves and bark	Ethanolic extract	Hydrolysable tannins, flavonoids and condens	In-vivo	-----	90% mortality rate at 100 ppm	

						ed tannins				
	Brazil	Eugenia dysenterica	Myrtaceae	Leaves and bark	Ethanolic extract	Condens ed tannins, flavanoi ds, phenolic acid and coumari ns	In-vivo	-----	100 % mortality rate	
	Brazil	Annona crassiflora	Annonaceae	Leaves	Ethanolic extract	Flavanoi ds and tannin	In-vivo	-----	Less than 10 % mortality rate	
	Puerto Rico	Didymopanax morototoni	Araliaceae	Leaves	Methanolic extract	----- --	In-vivo	-----	100 % mortality rate at 50 ppm	[121]
	Puerto Rico	Mammea americana	Clusiaceae	Leaves	Methanolic extract	----- --	In-vivo	-----	100 % mortality rate	
	Puerto Rico	Furcraea tuberosa	Agavaceae	Leaves	Methanolic extract	----- --	In-vivo	-----	100 % mortality rate	
	Puerto Rico	Argemone mexicana	Papaveracea e	Leaves	Methanolic extract	----- --	In-vivo	-----	67 % mortality rate at 100 ppm	
	Puerto Rico	Paullinia pinnata	Sapindaceae	Leaves	Methanolic extract	----- --	In-vivo	-----	50 % mortality rate at 50 ppm	
	Puerto Rico	Solanum americanum	Solanaceae	Leaves	Methanolic extract	----- --	In-vivo	-----	33 % mortality rate at 50 ppm	

	Brazil	Annona crassiflora	Annonaceae	Pulp, seed, stem and root	Ethanolic extract	----- --	In-vivo	LD90 (ppm): For adult= 2.34 For egg= 1	80 % mortality rate	[122]
	Brazil	Annona glabra	Annonaceae	Leaves and seed	Ethanolic extract	----- --	In-vivo	LD90 (ppm): For adult= 3.79 For egg= 1	80 % mortality rate	
	Brazil	Annona muricata	Annonaceae	Leaves and stem bark	Ethanolic extract	----- --	In-vivo	LD90 (ppm): For adult= 3.79 For egg= 1	60 % mortality rate	
	Brazil	Annona pisonis	Annonaceae	Leaves and stem bark	Ethanolic extract	----- --	In-vivo	LD90 (ppm): For adult= 6.21 For egg= 1	80 % mortality rate at 60ppm	
	Brazil	Annona salmani	Annonaceae	Leaves	Ethanolic extract	----- --	In-vivo	LD90 (ppm): For adult= 0.66	100 % mortality rate	

								For egg= 1		
	Brazil	Annona squamosa	Annonaceae	Seed, stem, root, and leaves	Ethanollic extract	----- --	In-vivo	LD90 (ppm): For adult= 8.55 For egg= 1	100 % mortality rate	
	Brazil	Euphorbia splendens	Euphorbiaceae	Latex	Aqueous solution	----- --	In-vivo	LD50= 0.60ppm LD90= 0.83 ppm	Extract proved toxic against snail sp.	[123]
	Iraq	Cucumis melo	Cucurbitaceae	Leaves and stem	Aquatic extract	----- --	In-vivo	LC50 (g/L): Leaves=40, Stem=50	15 % mortality rate with leaves extract while 12.5 % mortality rate with stem extract	[124]
	Iraq	Thymus Vulgaris	Lamiaceae	Leaves	Aqueous extract	----- --	In-vitro	LC50= 18.7 g/L	50 % mortality at 18.7g/L	[125]
	Iraq	Ricinus Communis	Euphorbiaceae	Leaves and bark	Aqueous extract	----- --	In-vivo	LC50 for 50 g/L Bark=191 % Leaves= 32%	Abnormal behavioral patterns	[126]
	France	Croton campestris	Euphorbiaceae	Root and bark	dichloromethanolic extract	velamone,	In-vivo	LD100 =50 ppm	100 % mortality rate	[127]

						velamolo ne and velamolo ne acetate				
	France	<i>Caryopteris clandonensis</i> x	Lamiaceae	Root and bark	Chloroform extract	Fuerstio ne, α- Caryopte rone and 15- deoxyfur estione	In-vivo	LD100= <3 ppm	100 % mortality rate	[128]
	United Kingdom	<i>Euphorbia myrsinites</i>	Euphorbiace ae	Leaves and stem	Aqueous extract	----- --	In-vivo	LC50 = 8.9 ppm	75 % reduced activity and 100 % mortality rate	[129]
	Belgium	<i>Maesa ceolata</i>	Myrsinaceae	Leaves	Aqueous extract	Saponin	In-vivo	-----	>70 % mortality rate	[130]
	Germany	<i>Jatropha curcas</i>	Euphorbiace ae	Seed	Methanolic extract	----- --	In-vitro	LC100 =25 ppm	100 % mortality rate	[131]
	Egypt	<i>Nigella sativa</i>	Ranunculace a	Seed	Oil	----- --	In-vitro	LC50= 5.8 ppm	Laceration of vacuolated tissues in foot region	[132]
	Egypt	<i>Pelargonium graveolens</i>	Geraniaceae	Seed	Oil	----- -	In-vitro	LC50= 15.0 ppm	Significant miracidicidal and cercaricidal activity	

	Egypt	<i>Azadirachta indica</i>	Meliaceae	Seed	Oil	----- -	In-vitro	LC50= 28.0 ppm	Formation of vacuoles in cytoplasm of digestive glands	
	Egypt	<i>Euphorbia pulcherima</i>	Euphorbiaceae	Whole plant	Methanolic extract	----- --	In-vivo	LC50= 43 ppm	After 3 hour, mortality rate of cercariae is 50 % at 15 ppm, 100 % mortality rate of miracidia at 100 ppm, 28.4 % reduction in total protein and 40 % decrease in lactate dehydrogenase	[133]
	Egypt	<i>Atriplex nummularia</i>	Amaranthaceae	Whole plant	Methanolic extract	----- --	In-vivo	LC50= 32 ppm	After 3 hour, mortality rate of cercariae is 70 % at 15 ppm, 100 % mortality rate of miracidia at 100 ppm, 51 % reduction in total protein and 62.1 % decrease in lactate dehydrogenase	
	Mali	<i>Glinus oppositifolius</i>	Aizoaceae	Whole plant	Crude extract	12- Glucosyl- GO1	In-vivo	LC50 of Butanol extract= 64.3 ppm, LC50 of Ethanolic extract= 86.2 ppm	100 % mortality rate of both snail species at 200 ppm	[68]
	Egypt	<i>Sesbania sesban</i>	Leguminosae	Leaves	Methanolic extract	----- --	In-vivo, in-vitro	LC50= 15 ppm	Reduced cercarial production and 44 % mortality after 48h	[134]

	Nigeria	Talinum triangulare	Talinaceae	Root	Ethanollic extracts	----- --	In-vivo	LC50= 251 ppm	Ovicidal activity at <300 ppm	[135]
	Egypt	Adenium obesum	Apocynaceae	Leaves	Methanolic extract	----- --	In-vivo	LC25=26.3 %	Reduced protein content in tissues	[136]
	Sudan	Cymbopogon nervatus	Poaceae	Leaves	Essential oils	----- --	In-vivo	LD50 = 237.33 ppm	100 % mortality rate at 400 ppm	[80]
	Sudan	Boswellia papyrifera	Burseraceae	Leaves	Essential oils	----- --	In-vivo	LD50= 311.05 ppm	90 % mortality rate at 350 ppm	
	Morocco	Withania somnifera	Solanaceae	Leaves and fruit	Aqueous and methanolic extract	Flavanooids, terpenes, alkaloids , saponins , tannins	In-vivo	LC50 = 94 ppm	Proved toxic to snail specie	[137]
	Morocco	Withania frutescens	Solanaceae	Root, leaves and fruit	Aqueous extract	Terpenes, alkaloids , tannins and saponins	In-vivo	-----	Proved toxic to snail specie	
	Morocco	Solanum elaeagnifolium	Solanaceae	Fruit, Root	Aqueous extract	Alkaloids , tannins and saponins	In-vivo	LC50 = 9.5 ppm	Potential molluscicide against snail	

	Morocco	<i>Silene cucubalus</i>	Caryophyllaceae	Root	Aqueous and n-butanol extract	Saponins, tannins and terpenes	In-vivo	-----	Potential molluscicide against snail	
	Morocco	<i>Spergularia marginata</i>	Caryophyllaceae	Root	Aqueous extract	Saponins	In-vivo	-----	Proved toxic to snail specie	
	Morocco	<i>Saponaria vaccaria</i>	Caryophyllaceae	Root	Aqueous extract	Saponins	In-vivo	-----	Potential molluscicide against snail	
	Sudan	<i>Azadirachta indica</i>	Meliaceae	Leaves and seeds	Aqueous extract	----- --	In-vivo	-----	75 % decrease in snail species after 5 days	[84]
	Egypt	<i>Commiphora molmol</i>	Bruseraceae	Whole plant	Oil extract	----- --	In-vivo	LD50= 50 ppm LD90= 95 ppm	100 % mortality rate	[53]
	Morocco	<i>Citrus aurantium</i> var. valencia	Rutaceae	Fruit	Essential oil	α -pinene, sabinene, myrcene and linionene	In-vivo	LC50= 0.28 ppm	Toxic against the snail sp.	[138]
	Morocco	<i>Origanum compactum</i>	Lamiaceae	Aerial parts	Essential oil	Thymol, α -terpineol, γ -terpinene	In-vivo	LC50= 0.44 ppm	Toxic against the snail sp.	

						e and α -pinene				
	Mali	Cussonia barteri	Araliaceae	Roots	Methanolic and aqueous extract	----- ---	In-vitro	-----	Extract proved toxic against snail sp.	[85]
	Mali	Glinus oppositifolius	Aizoaceae	Whole plant	Methanolic and aqueous extract	----- ---	In-vitro	-----	Extract proved toxic against snail sp.	
	Mali	Lannea velutina	Anacardiaceae	Leaves	Methanolic and aqueous extract	----- ---	In-vitro	-----	Extract proved toxic against snail sp.	
	Egypt	Commiphora molmol	Burseraceae	Whole plant	Oil and resin extract	----- ---	In-vivo	LC50= 3 ppm	100 % mortality rate	[56]
	Morocco	Ruta chalepensis L.	Rutaceae	Aerial part	Aqueous and ethanolic extract	Phenol, coumarins, flavanoid, alkaloid, sterol and terpenoids	In-vivo	LC50= 1.41 ppm LC90= 2.23 ppm	Strong molluscicidal activity	[139]
	Morocco	Zygophyllum gaetulum Emb.	Zygophyllaceae	Aerial part	Aqueous and ethanolic extract	Flavonoid and saponin	In-vivo	LC50= 8.69 ppm LC90= 16.82 ppm	High mortality rate	

	Morocco	Citrus bigaradia Duhamel	Rutaceae	Leaves	Aqueous and ethanolic extract	Terpenoid, phenol, coumarins, flavanoid, alkaloid, sterol and terpenoid	In-vivo	LC50= 2.74 ppm LC90= 4.43 ppm	Toxic for adult snails
	Morocco	Capparis spinosa L.	Capparidaceae	Aerial part	Aqueous and ethanolic extract	Coumarin, alkaloid and sterol	In-vivo	LC50= 70.58 ppm LC90= 122 ppm	Strong molluscicidal activity
	Morocco	Chenopodium ambrosioides L.	Chenopodiaceae	Aerial part	Aqueous and ethanolic extract	Flavonoid, saponin and terpenoid	In-vivo	LC50= 1.41 ppm LC90= 2.23 ppm	Strong molluscicidal activity
	Morocco	Artemisia herba alba Asso.	Asteraceae	Aerial part	Aqueous and ethanolic extract	Phenol, flavanoid, alkaloid, sterol and terpenoid	In-vivo	LC50= 62.94 ppm LC90= 92 ppm	Strong molluscicidal activity

	Morocco	<i>Origanum compactum</i>	Lamiaceae	Aerial part	Aqueous and ethanolic extract	Phenol, flavanoid, saponin and terpenoid	In-vivo	LC50= 8.22 ppm LC90= 1.99 ppm	Strong molluscicidal activity
	Morocco	<i>Melia azedarach</i> L.	Meliaceae	Leaves	Aqueous and ethanolic extract	Flavonoid, alkaloid and terpene	In-vivo	LC50= 3.53 ppm LC90= 4.20 ppm	High mortality rate
	Morocco	<i>Delphinium staphisagria</i> L.	Ranunculaceae	Flower	Aqueous and ethanolic extract	Alkaloid and saponin	In-vivo	LC50= 3.5 ppm LC90= 8.39 ppm	Toxic for adult snails
	Morocco	<i>Nigella sativa</i>	Ranunculaceae	Aerial part	Aqueous and ethanolic extract	Coumarins, alkaloid, terpenoid, saponin and sterol	In-vivo	LC50= 106.7 ppm LC90= 140 ppm	Toxic for adult snails
	Morocco	<i>Zizyphus vulgaris</i> Lamk.	Rhamnaceae	Leaves	Aqueous and ethanolic extract	Phenol, saponin and alkaloid	In-vivo	LC50= 4.12 ppm LC90= 8.41 ppm	High mortality rate

	Morocco	<i>Chrysanthemum viscidifolium</i>	Asteraceae	Aerial part	Aqueous and ethanolic extract	Flavonoid and terpenoid	In-vivo	LC50=4.43 ppm LC90=8.39 ppm	Strong molluscicidal activity	
	Morocco	<i>Citrullus colocynthis</i> L.	Cucurbitaceae	Fruit, Seed	Aqueous and ethanolic extract	Sterol and terpenoid	In-vivo	LC50= 4.35 ppm LC90= 8.39 ppm	Strong molluscicidal activity	
	Morocco	<i>Lavandula stoechas</i> L.	Lamiaceae	Aerial Part	Aqueous and ethanolic extract	Phenol, flavonoid and terpenoid	In-vivo	LC50= 106.7 ppm LC90= 140 ppm	High mortality rate	
	China	<i>Solanum xanthocarpum</i>	Solanaceae	Fruit	Ethanolic extract	----- --	In-vivo	LC50 (mg/l): =1.35	100 % mortality rate	
	Germany	<i>Jatropha curcas</i>	Euphorbiaceae	Seed	Methanolic extract	----- --	In-vitro	LC100= 1 ppm	100 % mortality rate of snail specie	[131]
<i>Bulinus globosus</i>	Zimbabwe	<i>Cucurbita maxima</i>	Cucurbitaceae	Seed	Aqueous and crude ethanolic extracts	----- --	In-vivo	LC50 (mg/ml) for: Aqueous extracts on <i>B. globosus</i> adults= 0.004	100 % mortality rate at 2 mg/mL	[59]

								Ethanollic extracts on B. globosus adults= 0.19		
	Nigeria	Allium cepa, Allium sativum	Amaryllidaceae	Bulb and cloves	Methanolic extract	----- ---	In-vivo	LC50 (60 mg/l): For Allium cepa=15; For Allium sativum= 19.37	Mortality is higher in case of Allium sativum than Allium sepa and highest mortality rate is observed in 42.15 mg/l	[140]
	Nigeria	Balanites aegyptiaca	Zygophyllaceae	Stem and bark	Aqueous solution	2-methoxy-4-vinylphenol, benzaldehyde, phenol, 2-methoxy-4-(2-propenyl)-acetate, ethanone, 1-(4hydroxyphenyl)-, alpha-	In-vivo	LD50 (mg/L): For juvenile= 5.7 For adult= 6.2 LD90 (mg/L): For juvenile= 6.8 For adult= 7.2	100 % mortality rate of both snail species at 40mg/L	[67]

						d-6,3-furanose, 4-(3-hydroxy-1-propenyl)-2-methoxy phenol, hexadecanoic acid, 2-butanone, 9-octadecanoic acid, octadecanoic acid, 13-octadecenal, tetradecanoic, 9-octadecanamide and di-n-octylphthalate				
	Nigeria	Rhizophora mangle	Rhizophoraceae	Leaves	Methanolic extract	Saponins, glycosides, tannins, alkaloid	In-vivo	LC50 = 87.50 ppm	90 % mortality rate at 200 ppm	[141]

						and flavanoids				
	Nigeria	Rhizophora racemosa	Rhizophoraceae	Leaves	Methanolic extract	Saponins, glycosides, tannins, alkaloid and flavanoids	In-vivo	LC50 = 125 ppm	100 % mortality rate at 250 ppm	
	Nigeria	Avicennia germinans	Verbenaceae	Leaves	Methanolic extract	Saponins, glycosides, tannins, alkaloid and flavanoids	In-vivo	LC50 = 87.50 ppm	80 % mortality rate at 150 ppm	
	Nigeria	Languncularia racemosa	Combretaceae	Leaves	Methanolic extract	Saponins, glycosides, tannins, alkaloid and flavanoids	In-vivo	LC50 = 102.09 ppm	100 % mortality rate at 50 ppm	

	Nigeria	<i>Azadirachta indica</i>	Meliaceae	Root and bark	Ethanol, chloroform, acetone and n-hexane extract	Triterpenes, flavanoids, glycosides and tannins	In-vivo	LC50 = 0.35 ppm	100 % mortality rate	[142]
	Nigeria	<i>Talinum triangulare</i>	Portulacaceae	Leaves and root	Ethanol extract	-----	In-vivo	LC50= 125.89 ppm	100 % mortality at 160 ppm (root) and 640 ppm (leaves)	[143]
	Nigeria	<i>Azadirachta indica</i>	Meliaceae	Leaves, seed and bark	Methanol and decoction extract	Saponin, tannin, flavanoid and glycoside	In-vitro	-----	100 % mortality rate after 24 h	[144]
	Nigeria	<i>Hyptis suaveolens</i>	Lamiaceae	Leaves	Ethanol extracts	----- --	In-vivo	At egg stage LC50= 0.614 ppm; At adult stage LC50= 0.077 ppm	80 % mortality rate at 0.8 ppm	[145]
	Nigeria	<i>Securidaca longepedunculata</i> , <i>Tephrosia brateolata</i>	Polygalaceae, Leguminosae	Root, leaves, stem and bark	Ethanol and methanol extract	----- --	In-vivo	With ethanolic extract: LC50 (ppm) leaf= 0.15 bark = 0.19 Root= 0.18	100 % mortality rate	[146]

								With methanolic extract: LC50 (ppm) leaf = 0.55 bark= 0.60 root= 0.21		
	Nigeria	Carica papaya	Caricaceae	Leaves	Ethanolic extract	----- --	In-vivo	LC50 = 619.1 ppm LC90= 1180.7 ppm	100 % mortality rate	[79]
	Nigeria	Terminalia catappa	Combretacea e	Leaves	Ethanolic extract	----- --	In-vivo	LC50 = 1095.7 ppm LC90 = 1874.9 ppm	100 % mortality rate	
Oncomelania hupensis	China	Camellia oleifera	Theaceae	Seed	Aqueous solution	Tea-seed distilled saponin	In-vivo	LD50 at 24 hr in laboratory = 0.701 mg/L	Mortality rate is 72-98 %	[147]
	China	Macleaya cordata	Papaveracea e	Fruit	Aqueous solution	Quatern ary benzo[c] phenant hridine	In-vivo, In-vitro	LC50=1.29 mg/L; LC90=2.92 mg/L	Decrease in glutathione stransferase	[148]

						alkaloids (sanguinarine and chelerythrine) and protopine alkaloids (protopine and α -allocryptopine)			(GST), carboxylesterase, acid phosphatase and alkaline phosphatase activity	
	Philippines	<i>Jatropha curcas</i> L.	Euphorbiaceae	Leaves	Petroleum ether extract	Alkaloids, tannins, terpenoids, flavonoids and saponins	In-vivo	LD50 = 0.45 mg/mL	80 % mortality at 1.5 mg/mL	[149]
	China	<i>Macleaya cordata</i> (Wild) R.	Papaveraceae	Leaves	Aqueous solution	Alkaloids	In-vivo	LC50 = 6.35 mg/L LC90 = 121.23 mg/L	Mortality rate is 73 % at 25 mg/L, Increase in alanine aminotransferase, aspartate transaminase and decrease in esterase isozyme	[150]

	China	Buddleja lindleyana	Loganiaceae	Whole plant	Ethanolic extract	Acacetin-7-rutinoside	In-vivo	LC50 = 36:12 mg/L	100 % mortality at 70 mg/L	[151]
	China	Cinnamomum camphora	Lauraceae	Leaves	-----	Linalool, α -terpineol, 4-terpineol, camphora, β -phellandrene, 1,8-cineole and spathulenol	In-vitro	LC50 = 0.25 mg/L	Disfigurement noted in hepatopancrease and gills	[152]
	China	Ginkgo biloba	Ginkgoaceae	Leaves and sarcotesta	Ethanolic extract	-----	In-vivo	LD50= 47.64 mg/L	Negatively affect the functioning of dihydronicotinamide adenine dinucleotide, cytochrome c oxidase, cytochrome b and (NADH) dehydrogenase, adenosine triphosphate (ATP) synthase	[153]
	China	Pulsatilla chinensis	Ranunculaceae	Whole plant	Alcoholic extract	Pulchelinonin A3 and	In-vitro	LC50 = 0.48 mg/L	Significant inhibition of cholinesterase, aspartate	[154]

						Pulchinelloside B			transaminase and alkaline phosphatase	
	China	<i>Sapium sebiferum</i>	Euphorbiaceae	Leaves, branch, root and seed	Aqueous, ethyl acetate and ethnolic extract	----- ---	In-vivo	-----	Decreased protein content in liver and 100 % mortality rate	[155]
	China	<i>Macleaya cordata</i>	Papaveraceae	Fruit	Dechlorinized aqueous extract	Bisulfates of sanguinarine, total alkaloids and chelerythrine	In-vivo	LC50= 0.19 mg/L	100 % mortality rate at 72h	[156]
	China	<i>Eomecon chionantha</i>	Papaveraceae	Root stalk	Alcoholic extract	Alkaloids	In-vivo	-----	100 % mortality rate at 2.5 mg/l and increased amount of alanine aminotransferase	[157]
	China	<i>Ginkgo biloba</i>	Ginkgoaceae	Leaves	Ethanollic and aqueous extract	----- --	In-vivo	-----	80 % mortality rate after 24 h, decreased total protein content and glycogen	[158]
	China	<i>Torreya grandis</i>	Taxaceae	Leaves	Ethanollic and aqueous extract	----- --	In-vivo	-----	72 % mortality rate after 72 h	

	China	Ailanthus altissima	Simaroubaceae	Leaves	Ethanolic and aqueous extract	----- --	In-vivo	-----	40 % mortality rate after 72 h	
	China	Peucedanum praerutorum	Apiaceae	Root	Ethanolic and aqueous extract	----- --	In-vivo	-----	83 % mortality rate after 72 h	
	China	Herba agrimoniae	Rosaceae	Whole plant	Ethanolic and aqueous extract	----- --	In-vivo	-----	80 % mortality rate after 72 h	
	China	Buddleja lindleyana	Scrophulariaceae	Leaves	Ethanolic extract	----- --	In-vivo	LC50 = 39.91 mg/L LC90 = 59.28 mg/L	96 % mortality rate after 48 h	[159]
	China	Eupatorium adenophorum	Asteraceae	Leaves, stem and root	Aqueous extracts	----- --	In-vivo	-----	100 % mortality rate after 58 h	[160]
	China	Ginkgo biloba	Ginkgoaceae	Seed	Petroleum ether extracts	Ginkgolide acids	In-vivo	LD50 = 7.81ppm	100 % mortality rate after 48h	[161]
	China	Arisaema erubescens	Araceae	Tubers	Ethanolic extract	----- --	In-vitro	LC50= 60.2 mg/L LC90= 145.6 mg/L	Increased activity of <i>alkaline phosphatase and choline esterase</i>	[162]
	China	Ginkgo biloba	Ginkgoaceae	Seed	Benzinum extracts	----- --	In-vivo	LC50= 0.65 mg/L	Decreased activity of alkaline phosphatase, choline esterase,	[163]

								LC90= 5.5 mg/L	malic dehydrogenase and alanine aminotransferase	
	China	<i>Solanum xanthocarpum</i>	Solanaceae	Fruit	Ethanolic extract	----- --	In-vivo	LC50 = 0.62 mg/l	100 % mortality rate	[164]
Bulinus rholfsi	Nigeria	<i>Rhizophora mangle</i>	Rhizophorac eae	Leaves	Methanolic extract	Saponins , glycosid es, tannins, alkaloid and flavanoi ds	In-vivo	LC50 = 108.22 ppm	90 % mortality rate at 200 ppm	[141]
	Nigeria	<i>Rhizophora racemosa</i>	Rhizophorac eae	Leaves	Methanolic extract	Saponins , glycosid es, tannins, alkaloid and flavanoi ds	In-vivo	LC50= 83.51 ppm	100 % mortality rate at 250 ppm	
	Nigeria	<i>Avicennia germinans</i>	Verbenaceae	Leaves	Methanolic extract	Saponins , glycosid es, tannins, alkaloid and	In-vivo	LC50 = 123.74 ppm	80 % mortality rate at 150 ppm	

						flavanoi ds				
	Nigeria	Languncularia racemosa	Combretacea e	Leaves	Methanolic extract	Saponins , glycosid es, tannins, alkaloid and flavanoi ds	In-vivo	LC50= 152.03 ppm	100 % mortality rate at 50 ppm	
Bulinus natalensis	Germany	Jatropha curcas	Euphorbiace ae	Seed	Methanolic extract	----- ---	In-vitro	LC100= 1 ppm	100 % mortality rate of snail specie	[165]
Lymnaea luteola	India	Sapindus trifoliatus	Sapindaceae	fruit	Methanolic extract	----- ---	In-vitro	LC50= 50 ppm LC90= 87 ppm	80 % mortality rate	[166]
	India	Acacia concinna	Leguminosa e	fruit	Methanolic extract	----- ---	In-vitro	LC50= 360 ppm LC90= 451 ppm	90 % mortality rate at 300 ppm	
	India	Madhuca indica	Sapotaceae	Seed	Methanolic extract	----- ---	In-vitro	LC50= 62 ppm LC90= 100 ppm	90 % mortality rate at 80 ppm	

	India	Phytolacca acinos	Phytolaccaceae	fruit	Methanolic extract	----- ---	In-vitro	LC50= 193 ppm LC90= 399 ppm	80 % mortality rate	
Monacha cartusiana	Egypt	Syzygium aromaticum	Myrtaceae	Flower bud	Acetone and ethanol extract	----- -	In-vivo	LC50 =1.06; LC95=26.8 3	Excessive mucosal secretion, hyperactivity and 40-60 % mortality rate	[167]
	Egypt	Zingiber officinale	Zingiberaceae	Rhizome	Ethanolic extract	----- ---	In-vitro, In-vivo	-----	90 % mortality at 28th day, Increased cell lysis and vacuole formation in digestive gland with disrupted calcium spherules and accumulation of pyknotic nuclei, increased levels of ALT, AST, α & β esterases and phenoloxidas, deformed egg and sperm	[168]
	Egypt	Balanites aegyptiaca	Zygophyllaceae	Fruit	Ethanolic extract	Diosgenin and 3,5- spirostadiene	In-vivo	LC50= 0.256 ppm	73 % mortality rate	[169]
Monacha Obstructa	Egypt	Ricinus communis	Euphorbiaceae	Fruit and leaves	Aqueous and ethanolic extract	Ricinisine, ricinoleic acid, linoleic acid and quinic acid	In-vivo	LC50= 0.215 LC90= 16.128	80 % mortality rate after 72 hr	[170]

	Egypt	Anagallis arvensis L.	Primulaceae	Leaves and fruit	Hexane extract	Malonic acid, aromatic acid, 2-tertbutyl-4-isopropyl-5-methylphenol and phosphoric acid tributyl ester	In-vivo	LC50 after 24h= 1.70 g/ml	Extract induced 34 % toxicity to adult snails	[171]
	Egypt	Solanum nigrum L.	Solanaceae	Leaves and fruit	Hexane extract	-----	In-vivo	LC50 after 24h= 2.75 g/ml	Proved 75 % toxic	
	Egypt	Cyperus longus L.	Cyperaceae	Rhizome	Hexane extract	-----	In-vivo	LC50 after 24h= 3.09 g/ml	Acted as bait (24.2 %) after 96 h	
	Saudi Arabia	Adenium obesum	Apocynaceae	Stem	Hydro-ethanolic extract	Cerberin and neriifolin	In-vivo	LD50 (µg/g): Of Cerberin = 4.3 Of neriifolin = 5.39	Toxicity index of Cerberin is 79.81 while of neriifolin is 100	[172]

Indoplanorbis exustus	India	Artemisia annua	Asteraceae	Leaves	Crude extract	----- -	In-vitro	LC50= 5.9 ppm	Abnormal behavior (rapid movement)	[173]
	India	Leucas aspera, Parthenium hysterophorus	Lamiaceae, Asteraceae	Leaves	Aqueous solution	----- ----	In-vivo	-----	90 % mortality rate	[174]
Monacha cantiana	Saudi Arabia	Calotropis procera, Adenium arabicum	Apocynaceae	Latex and stem	Cardenolide and methonyl extract	Glycosides	In-vivo	LD50 values 12.62 mg/kg for cardenolide LD50 for methomyl was 116.62 mg/kg	80 % mortality rate	[175]
Lanistes ovum	Nigeria	Senna occidentalis	Leguminosae	Leaves	Ethanolic extract	Tannins, inulin, flavanoids and alkaloids	In-vitro	LC50=83.67 mg/L	100 % mortality rate after 96h	[176]
	Nigeria	Vernonia amygdalina	Asteraceae	Leaves	Ethanolic extract	Tannins, alkaloids, flavanoids and saponin	In-vitro	LC50=614.288 mg/L	100 % mortality rate after 96h	

	Nigeria	Khaya senegalensis	Meliaceae	Leaves	Ethanollic extract	Tannins, flavanoids, alkaloids, saponins and inulin	In-vitro	LC50=395.58 mg/L	Inhibition of alpha-glucosidase and reduced glucose level	
Physa acuta	Iraq	Melia azadirachta	Meliaceae	Fruit	Fruit extract	----- -	In-vivo	At 24th hr, LC50=10 ppm ; LC90 4.4 ppm	Reduced juvenile movement, Irregular heartbeat and reduced to 58 beats/min from 88 beats/min, 100 % mortality rate at 5 ppm	[177]
Cornu aspersum	Lebanon	Styrax officinalis	Styracaceae	Fruit	Ethanollic extract	Saponins, triterpenes and tannins	In-vivo	-----	Disintegrated membrane tissues	[178]
Subulina octona	Brazil	Mikania glomerata	Asteraceae	Leaves	Aqueous extract	Tannins	In-vivo	LC50=45 mg/mL	100% mortality rate at 60th day	[179]
	Brazil	Bidens pilosa Linné	Asteraceae	Aerial parts	Aqueous extract	Flavonoids and saponin	In-vivo	LC50=51.4 mg/mL; LC90=74.1 mg/mL	Reduced hatchability to 61 %, 100 % mortality rate after 48 h	[180]
Lanistes lybicus	Nigeria	Erythrophleum suaveolens	Leguminosae	Stem and bark	Ethanollic extract	Saponin	In-vivo	38.74 µg/ml	100 % mortality rate at 50 µg/ml	[181]
Physa marmorata	Brazil	Moringa oleifera	Moringaceae	Seed	Aqueous extract	----- -	In-vivo	P. marmorata	Toxic for Physa marmorata	[107]

								LC50 =0.339 g/l; LC90 =0.789 g/l		
Bradybaena similaris	Brazil	Solanum paniculatum	Solanaceae	Leaves	Aqueous extract	----- -	In-vivo	-----	75 % mortality rate after 24 h, significant reduction of glucose, glycogen and galactogen	[182]
Bulinus sp.	Kenya	Balanites aegyptiaca	Balanitaceae	Leaves, root, stem and fruit	Methanolic extract	----- -	In-vitro	LD50= 2.0931	38 % mortality rate	[183]
	Kenya	Phytolacca dodecandra	Phytolaccaceae	Leaves, root, stem and fruit	Methanolic extract	----- -	In-vitro	LD50= 2.5332	32 % mortality rate	
	Kenya	Phytolacca octandra	Phytolaccaceae	Leaves, root, stem and fruit	Methanolic extract	----- -	In-vitro	LD50= 2.2627	36 % mortality rate	
	Nigeria	Moringa oleifera	Moringaceae	Seeds	Aqueous extract	----- -	In-vivo	-----	100 % mortality rate at 2000 ppm	[184]
Biomphalaria juvenile snails and Bulinus adult snails	Zimbabwe	Curcubita maxima	Cucurbitaceae	Seed	Aqueous extract	----- -	In-vivo	LC50 (mg/ml): On Biomphalaria juvenile= 0.002	100 % mortality rate	[185]

								On Bulinus adult = 0.004		
Different snail sp.	Sudan	Khaya grandifoliola	Meliaceae	Bark	Aqueous and ethanolic extract	----- -	In-vivo	-----	100 % mortality rate at 1 g/L	[186]

A Most of the plants exhibiting molluscicidal activity belonged to the families Euphorbiaceae (9.31 %); Solanaceae (7.16 %); Asteraceae (6.09 %); Leguminosae (5.73 %); Lamiaceae (4.3 %); Meliaceae (3.94 %); Myrtaceae and Annonaceae (2.86 %); Rutaceae and

Zygophyllaceae (2.5 %); Moringaceae, Combretaceae, and Araliaceae (2.15 %), as presented in Figure 2. A number of other families (n= <5; <1.79 %; Table 1) also included plants showing efficacy against various snail species.




Figure 2: Major plant families exhibiting molluscicidal activity against intermediate snail hosts of schistosomes.

As shown in Figure 3, the key plant parts used to assess molluscicidal activity included leaves (48.74 %), fruit (15.05 %), root (14.69 %), stem (12.54 %), seed (12.9 %), bark (10 %), whole plant (9.31 %), and flower (2.15 %).




Figure 3: Pie chart depicting the percentage of plant parts used to assess molluscicidal activity.

Discussion

Plant-based molluscicides are receiving increasing recognition because of their higher biodegradability and low cost [63]. These molluscicides include plant secondary metabolites belonging to the category's tannins, glycosides, terpenoids, alkaloids, and phenols. Some of the important phytoconstituents are described below. Saponins are chemical compounds belonging to the category of glycosides. They are further divided into monodesmosidic and bidesmosidic saponins and are effective molluscicides because of their toxicity. One of the studies [18] indicated that the killing effect of *Phytolacca dodecandra* on snail species was attributed to saponins. In another study [187], saponins present in *Ocimum americanum* increased the mortality rate of *Biomphalaria pfeifferi*. Moreover, bidesmosidic saponins extracted from *Yucca desmettiana* leaves effectively reduced the occurrence of *Biomphalaria alexandrina*; the lethal concentration (LC₅₀) after 24 h of exposure was 68 mg/L [41].

The concept of essential oils (EOs) is increasingly gaining popularity since it has the potential to perform a wide range of beneficial functions such as combating bacteria, alleviating inflammation, and killing parasites, in particular. Recent research indicates that *Schistosoma* worms can be prevented to grow by using oils of various plants. Indicatively, A lipid extract of *Plectranthus neochilus* is very potent in worm-killing, which is primarily due to the presence of 2-caryophyllene and 2-pinene [188]. Likewise, oil of *Dysphania ambrosioides* is able to kill adult *Schistosoma mansoni* worms when placed in a laboratory

[189]. This implies that we must continue to screen EOs to identify additional plant chemicals effective against schistosomiasis. Similar effects have been found with other EOs, e.g. *Mentha x villosa* and *Foeniculum vulgare*, which is why it is believed that EOs may be natural remedies [190]. Analyzing EOs derived from plants can inform not only whether they are effective but also how they are: by rupturing the walls of cells of the parasite or by disrupting its most vital processes (Islam et al., 2020).

A study by [24], showed that alkaloids obtained from the leaf ethanolic extract of *Ziziphus spina-christi* led to a high apoptotic rate in *Biomphalaria alexandrina*. These alkaloids are organic compounds that are greatly soluble in chloroform and ethanol than water. They are present in every plant part and are proved highly toxic to the snails. Another study of Tripathi and Singh 2000, reported that isopelletierine isolated from *Punica granatum* significantly reduced the activity of *Lymnaea acuminata*. Alkaloids (sanguinarine) purified from *Macleaya cordata* also significantly decreased the activity of glutathione-S-transferase, carboxylesterase, and alkaline phosphatase in *Oncomelania hupensis* [150]. Flavonoids are flavone-based compounds in plants proven to exhibit molluscicidal activity. According to one of the studies [192], a well-known flavonoid (quercetin) isolated from the leaf ethanolic extract of *Polygonum senegalense* showed significant molluscicidal activity in *Biomphalaria pfeifferi* and *Biomphalaria glabrata* within 1 day of exposure. Flavonoids isolated from *Stryphnodendron adstringens* caused 90 % mortality in *Biomphalaria glabrata* [120]. Flavonoids are toxic for snail species, but flavone glycosides isolated from *Asparagus plumosus* were non-toxic [193].

Tannins are plant-based organic compounds divided into two groups: hydro-stable tannins and condensed tannins. Various plant species, including *Croton megalocarpus*, *Aloe secundiflora*, *Moringa oleifera*, *Ipomea batatas*, *Chromolaena odorata*, *Ocimum americanum*, *Stryphnodendron adstringens* [107,117,187,194] are reported to contain tannins that are highly toxic to snails. Tannins present in *Persea Americana* enhanced the mortality rate of *Biomphalaria glabrata* embryos and caused lethargy in *Biomphalaria glabrata* adults [97]. Several plant species have been demonstrated to be effective against various snail species, but there are no sufficient data to determine their mechanism of action [195]. In 1998, WHO presented a report that emphasized the need to investigate the mode of action of molluscicidal plants, as they are economical, sustainable to use, and commonly available [196].

Plant molluscicides are classified on the basis of their mode of action into the following categories: enzyme inhibitors, neurotransmitter inhibitors (neurotoxins), stomach poisons, respiratory poisons, contact poisons, and growth inhibitors. The findings of some studies that investigated the mechanism of action of molluscicidal plants are discussed below: In recent years, a few studies have investigated the system of mollusks affected by molluscicides. [92] demonstrated that the latex of *Euphorbia cauducifolia* disrupted water absorption in *Biomphalaria glabrata*. The snail species that act as intermediate hosts of schistosomes are mainly in fresh water; therefore, the hemolymph of these snails is hyper-osmotic in nature. Consequently, it is crucial that the nutrients and water are in continuous equilibrium; their imbalance leads to reduced functioning and altered behavior (rapid and disoriented movement). It also causes tissue swelling, which makes snails more susceptible to plant molluscicides [8].

Plant molluscicides cause significant hormonal changes in snails, leading to a high mortality rate [14]. Enzymes such as acetylcholinesterase, succinate dehydrogenase, and alkaline phosphatase are used as indicators of plant molluscicidal activity. *Biomphalaria alexandrina* exposed to dry powdered *Datura stramonium* exhibited an increase in the levels of alanine transaminase and alkaline phosphatases [172]. Another study [49] showed that the methanolic extract of *Guayacum officinalis* decreased the levels of acetylcholinesterase and succinate dehydrogenase in *Biomphalaria alexandrina*. An aqueous extract of *Arisaema erubescens* was also shown to decrease the levels of superoxide dismutase and peroxidase in the liver of *Oncomelania hupensis*; in contrast, the malondialdehyde content increased, which indicates the oxidative stress

damage caused by the plant extract, leading to snail mortality [162]. Molluscicides may also act by negatively affecting the growth and development of snails. The findings of Ibrahim et al. [20] indicated that cerium oxide nanocomposites isolated from *Moringa oleifera* significantly reduced the reproductive rate of *Biomphalaria alexandrina*, with an LC₅₀ of 314.5 mg/L. The results of this study also indicated the deformation of hermaphrodite and digestive glands in the *Biomphalaria alexandrina*. Certain flavonoids are highly toxic to schistosomes and others are not. This disparity is based on their structure, which influences the interaction with cell and their availability within the body. The hydroxyl groups of flavonoid compounds tend to make the flavonoids better antioxidants and antiparasites when they are present compared to when they are absent [197]. Hence, the research on the structure-activity relationship (SAR) of these compounds should be examined by scientists to inform future studies on flavonoid drugs against schistosomiasis.

Schistosomiasis also depends on the temperature of the surrounding environment to be spread. Research demonstrates that increased temperatures accelerate *Bulinus* snail life cycle and lead to more cercariae being released, and other research illustrates that high temperatures cause stress in the snails and lower their parasite-carrying capacity [198]. Such mixed evidence implies we require more ecological models to anticipate how schistosomiasis will evolve with different temperatures, particularly with the rising heat extremes with climate change.

When snails encounter a toxic substance, they produce more mucus in an attempt to excrete the noxious substance and detoxify their body [196]. The mucus forms a protective barrier around the skin and reduces the harmful effect of molluscicides [199]. Similarly, several other factors have been reported to decrease the molluscicidal activity of plants. The surface-to-volume ratio of snails is another factor responsible for the variation in the efficacy of plant molluscicides. According to [59], the effect of *Cucurbita maxima* extracts on *Biomphalaria pfeifferi* juveniles and adults are different. The LC₅₀ values for adults and juveniles were 0.002 and 0.004 mg/mL, respectively. The reason is the less absorption of extract leads to more adult snail susceptibility towards toxic effect of molluscicides. Apart from these biological factors, certain chemical factors influence the susceptibility of snails to these plant extracts. For example, enzymatic analysis suggests that animals—more specifically, invertebrates—have mixed-function oxidases in their hepatopancreas that oxidize toxic chemical compounds to make them non-toxic [199].

Environmental factors such as light, temperature and extraction technique also affect the efficacy of plant-based molluscicides. A study by [112] indicated that an increase in temperature decreased the molluscicidal activity of *Euphorbia milli*. In contrast, an increase in temperature during phytochemical extraction helps these chemicals to easily penetrate the snail tissue [200]. Notably, [195] reported that the mortality rate of *Lymnea acuminata* treated with oleoresin decreased during winter.

Conclusion

This review highlights the significant potential of plant-derived compounds as eco-friendly molluscicides for controlling schistosomiasis. The evidence indicates that several plants and their phytochemicals exhibit strong molluscicidal activity, providing a promising alternative to synthetic chemicals. However, most studies remain limited to laboratory settings, and translation into large-scale field application is still scarce. Future research should focus on standardizing extraction methods, evaluating long-term safety and environmental impacts, and conducting field trials to validate laboratory findings. Collaborative efforts between researchers, policymakers, and public health authorities are essential to integrate plant-based molluscicides into schistosomiasis control programs. Such an approach could help reduce transmission while promoting sustainable and cost-effective disease management strategies.

Author Contributions: K.Q. collected the data and wrote the manuscript following discussions with N.A.S, H.A., and J.C. H.A. designed the study. K.Q. M.K performed the study. H.A., N.A.S., J.Z., and J.C. revised the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding

N/A

Informed Consent Statement

Not applicable

Data Availability Statement

The data presented in this study will be made available upon request the corresponding author.

Acknowledgments

We would like to thank all participants involved in this survey.

Conflicts of Interest

The authors declare no conflicts of interest.

References

1. Nelwan ML. Schistosomiasis: Life Cycle, Diagnosis, and Control. *Curr Ther Res Clin Exp* 2019;91:5–9. <https://doi.org/10.1016/j.curtheres.2019.06.001>.
2. Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. *The Lancet* 2014;383:2253–64. [https://doi.org/10.1016/S0140-6736\(13\)61949-2](https://doi.org/10.1016/S0140-6736(13)61949-2).
3. Qi YX, Huang MR, Sun HY, Wu XY, Liu ZT, Lu DB. Prevalence of depressive symptoms in patients with advanced schistosomiasis in China: A systematic review and meta-analysis. *JournalsPlosOrg* 2024;18. <https://doi.org/10.1371/JOURNAL.PNTD.0012003>.
4. Muhsin MA, Wang X, Kabole F, Zilabumba J, Yang K. The Indispensability of Snail Control for Accelerating Schistosomiasis Elimination: Evidence From Zanzibar. *Trop Med Infect Dis* 2022. <https://doi.org/10.3390/tropicalmed7110347>.
5. M. Trienekens SC, Faust CL, Besigye F, Pickering L, Tukahebwa EM, Seeley J, et al. Variation in Water Contact Behaviour and Risk of *Schistosoma Mansoni* (Re)infection Among Ugandan School-Aged Children in an Area With Persistent High Endemicity. *Parasit Vectors* 2022. <https://doi.org/10.1186/s13071-021-05121-6>.
6. Gong Y, Feng J, Luo Z-W, Xue J-B, Guo Z-Y, Zhang L, et al. Spatio-Temporal Heterogeneity of Schistosomiasis in China Based on Multi-Stage, Continuous Downscaling of Sentinel Monitoring 2021. <https://doi.org/10.21203/rs.3.rs-776529/v1>.
7. Barbosa FS, Pereira da Costa D, Arruda F. New field observations on the competitive displacement between two species of planorbid snails inhabiting northeastern Brazil. *Mem Inst Oswaldo Cruz* 1981;76:361–6. <https://doi.org/10.1590/S0074-02761981000400004>.
8. Allan F, Ame SM, Tian-Bi YNT, Hofkin B V., Webster BL, Diakit  NR, et al. Snail-related contributions from the schistosomiasis consortium for operational research and evaluation program including xenomonitoring, focal mollusciciding, biological control, and modeling. *American Journal of Tropical Medicine and Hygiene* 2020;103:66–79. <https://doi.org/10.4269/ajtmh.19-0831>.
9. Huot C, Clerissi C, Gourbal B, Galinier R, Duval D, Toulza E. Schistosomiasis Vector Snails and Their Microbiota Display a Phyllosymbiosis Pattern. *Front Microbiol* 2020;10:1–10. <https://doi.org/10.3389/fmicb.2019.03092>.
10. Research MN-CT, 2019 undefined. Schistosomiasis: life cycle, diagnosis, and control. Elsevier 2019;91:5–9. <https://doi.org/10.1016/J.CURTHERES.2019.06.001>.

11. Augusto R de C, de Mello-Silva CCC. Phytochemical molluscicides and schistosomiasis: What we know and what we still need to learn. *Vet Sci* 2018;5:1-9. <https://doi.org/10.3390/vetsci5040094>.
12. World Health Organization. Progress report 2001-2011 and strategic plan 2012-2020. 2010.
13. Inobaya M, Olveda R, Chau T, Olveda D, Ross A. Prevention and control of schistosomiasis: a current perspective. *Res Rep Trop Med* 2014;5:65-75. <https://doi.org/10.2147/rrtm.s44274>.
14. Zheng L, Deng L, Zhong Y, Wang Y, Guo W, Fan X. Molluscicides against the snail-intermediate host of *Schistosoma*: a review. *Parasitol Res* 2021;120:3355-93. <https://doi.org/10.1007/s00436-021-07288-4>.
15. Castle GD, Mills GA, Gravell A, Jones L, Townsend I, Cameron DG, et al. Review of the molluscicide metaldehyde in the environment. *Environ Sci (Camb)* 2017;3:415-28. <https://doi.org/10.1039/c7ew00039a>.
16. He P, Wang W, Sanogo B, Zeng X, Sun X, Lv Z, et al. Molluscicidal activity and mechanism of toxicity of a novel salicylanilide ester derivative against *Biomphalaria* species. *Parasit Vectors* 2017;10:1-11. <https://doi.org/10.1186/s13071-017-2313-3>.
17. Kashyap S, Khagta S, Guleria K, Arya V. Plants as Molluscicides: A recent update. *Int J Bot Stud* 2019;4:50-6.
18. Lemma A. Laboratory and field evaluation of the molluscicidal properties of *Phytolacca dodecandra*. *Bull World Health Organ* 1970;42:597-612.
19. Kloos H, McCullough FS. Plant Molluscicides. *J Med Plant Res* 1982;46:195-209.
20. Ibrahim AM, Mohamed F, Al-Quraishy S, Abdel-Baki AAS, Abdel-Tawab H. Green synthesis of Cerium oxide / *Moringa oleifera* seed extract nano-composite and its molluscicidal activities against *biomphalaria alexandrina*. *J King Saud Univ Sci* 2021;33:101368. <https://doi.org/10.1016/j.jksus.2021.101368>.
21. Ibrahim AM, Saleh HA, Zayed KM. *Colchicum Ritchii* flower: a new molluscicidal plant for *Biomphalaria alexandrina* snails and the infective stages of *Schistosoma mansoni alexandrina* snails and the infective stages of *Schistosoma mansoni*. *Molluscan Res* 2021;41:289-97. <https://doi.org/10.1080/13235818.2021.2003982>.
22. Ibrahim AM, Sayed SSM. Assessment of the molluscicidal activity of the methanolic seed extracts of *Ziziphus spina-christi* and *Carica papaya* on immunological and molecular aspects of *Biomphalaria alexandrina* snails. *Aquac Res* 2021;52:2014-24. <https://doi.org/10.1111/are.15050>.
23. Abou El-Nour MF. Evaluation of molluscicidal, miracidicidal and cercaricidal activities of crude aqueous extracts of *origanum majorana*, *ziziphus spina-christi* and *salvia fruticosa* on *schistosoma mansoni* and *schistosoma haematobium*. *Egypt J Aquat Biol Fish* 2021;25:913-33. <https://doi.org/10.21608/ejabf.2021.173661>.
24. Ibrahim AM, Ghoname SI, Mansour SM, El-Dafrawy SM. Effect of some medicinal plant extracts as molluscicidal and apoptotic agents on *biomphalaria alexandrina* snails. *Egypt J Aquat Biol Fish* 2020;24:291-300. <https://doi.org/10.21608/ejabf.2020.80284>.
25. Rabia Ibrahim AB, Gouda MSO, Elsayed H, Attiyah S. Efficiency of Three Extracts of *Carica papaya* as Molluscicidal and Anti- schistosomal Agents against *Biomphalaria alexandrina* and *Schistosoma mansoni* by Flow Cytometry Efficiency of Three Extracts of *Carica papaya* as Molluscicidal and Anti-schistosomal Ag. *J Pharm Res Int* 2020;32:31-41. <https://doi.org/10.9734/JPRI/2020/v32i1130577>.
26. A. Saad AH, Azzam AM, Mostafa BB, El-said KM, Hanafy RA. Improvement molluscicidal activity of *Anagallis arvensis* extracted by copper oxide nanoparticles against *Biomphalaria alexandrina* snails Abdel. *Egypt J Aquat Biol Fish* 2019;23:27-41.
27. El-Khayat HMM, Metwally KM, Abououf NA. Histopathological and Ultrastructural Studies on *Biomphalaria alexandrina* Snails Infected with *Schistosoma Mansoni* Miracidia and Treated with Plant Extracts. *Egypt J Hosp Med* 2018;71:2792-804. <https://doi.org/10.12816/0045846>.
28. Ibrahim AM, Ghoname SI. Experimental Parasitology Molluscicidal impacts of *Anagallis arvensis* aqueous extract on biological , hormonal , histological and molecular aspects of *Biomphalaria alexandrina* snails. *Exp Parasitol* 2018;192:36-41. <https://doi.org/10.1016/j.exppara.2018.07.014>.
29. Bahgat DM, Mossalem HS, Al-sayed E, Eldahshan OA, Singab ANB, Abu HM, et al. Influence of saponin fraction from *Albizia anthelmintica* on *Biomphalaria alexandrina* snail; the intermediate host of *Schistosoma mansoni* in Egypt. *Egypt J Aquat Biol Fish* 2018;22:231-40.
30. Mohamed AH, Osman GY, El-Eman MA, Abdel-Hamid H, E.M. Ali R. Deteriorations in biological and biochemical aspects of *Biomphalaria alexandrina* snails exposed to methanol extract of *Callistemon citrinus* leaves. *J Egypt Soc Parasitol* 2018;48:343-55.
31. El-Emam M, GY O, Abdel-Hamid H, Mohamed AH, R E M A. Determination of egg laying capacity, sex hormones and mortality of *Biomphalaria alexandrina* snails exposed to methanol extract from curcumin and the plants *Callistemon citrinus* and *Zingiber officinale*. *J Biosci Appl Res* 2017;3:97-109.
32. El-deeb FAA, Marie MS, Hasheesh WS, Sayed SSM. Factors affecting the molluscicidal activity of *Asparagus densiflorus* and *Oreopanax guatemalensis* plants and *Difenoconazole* fungicide on *Biomphalaria alexandrina* snails. *Comp Clin Path* 2016;1-9. <https://doi.org/10.1007/s00580-016-2263-8>.
33. Ghaly NS, Mina SA, Younis N. Schistosomicidal and molluscicidal activities of two Junipers species cultivated in Egypt and the chemical composition of their essential oils. *Journal of Medicinal Plants Research* 2016;10:47-53. <https://doi.org/10.5897/jmpr2015.5993>.
34. Hamed M, El-Amin S, Abdel-Fattah AS. Molluscicidal activity of some constituents isolated from

- Cestrum purpureum. Pharmacology Online 2015;2:59–72.
35. Ibrahim AM, Abdel-Gawad MM, El-Nahas HA, Osman NS. Studies on the Molluscicidal Activity of Agave Angustifolia and Pittosporum tobira on Schistosomiasis transmitting snails. Journal of Egyptian Society of Parasitology 2015;45:133–41.
36. El-sayed MM, El-nahas HA, El-Wakil EA. Molluscicidal Properties and Chemical Constituents of Euphorbia peploides. Australian Journal of Basic and Applied Sciences 2014;8:541–8.
37. Gohary E, Laila RA, Marwa GAM. Biochemical Effect of Three Molluscicide Baits Against the Two Land Snails, Monacha cantiana and Eobania vermiculata (Gastropoda: Helicidae). International Journal of Agricultural Research 2011;6:682–90.
38. Al-Sayed E, Hamid HA, Abu El Einin HM. Molluscicidal and antischistosomal activities of methanol extracts and isolated compounds from Eucalyptus globulus and Melaleuca styphelioides. Pharm Biol 2014;52:698–705. <https://doi.org/10.3109/13880209.2013.865240>.
39. Sheir SK, Mohamed AH, Osman GY, El-nabi SEH, Allam A. Delta Journal of Science Efficiency of Citrus limon extract on biological and molecular activities of Biomphalaria alexandrina snails. Delta Journal of Science 2013;36:186–200.
40. Abdel-haleem AA. Comparative karyological studies on the two Egyptian schistosome vectors , Biomphalaria glabrata and B . alexandrina , with reference to chromosomal aberrations due to Za ' ater plant. International Journal of Academic and Scientific Research 2013;1:1–7.
41. Diab Y, Ioannou E, Emam A, Vagias C, Roussis V. Desmettianosides A and B , bisdesmosidic furostanol saponins with molluscicidal activity from Yucca desmettiana. Steroids 2012;77:686–90. <https://doi.org/10.1016/j.steroids.2012.02.014>.
42. Rizk MZ, Metwally NS, Hamed MA, Mohamed AM. Experimental Parasitology Correlation between steroid sex hormones , egg laying capacity and cercarial shedding in Biomphalaria alexandrina snails after treatment with Haplophyllum tuberculatum. Exp Parasitol 2012;132:171–9. <https://doi.org/10.1016/j.exppara.2012.06.011>.
43. Mahmoud MB, Ibrahim WL, Abou-el-nour BM, El-emam MA, Youssef AA. Biological and biochemical parameters of Biomphalaria alexandrina snails exposed to the plants Datura stramonium and Sesbania sesban as water suspensions of their dry powder. Pestic Biochem Physiol 2011;99:96–104. <https://doi.org/10.1016/j.pestbp.2010.11.005>.
44. Bakry FA, Mohamed RT, El-hommosany K. Biological and biochemical responses of Biomphalaria alexandrina to some extracts of the plants Solanum siniacum and Artemisia judaica L . Pestic Biochem Physiol 2011;99:174–80. <https://doi.org/10.1016/j.pestbp.2010.12.001>.
45. El-din AS, El-sayed K, Mahmoud M. Effect of ethanolic extract of Dalbergia sissoo plant parts on Biomphalaria alexandrina snail , the intermediate host of Schistosoma mansoni. Journal of Evolutionary Biology Research Vol 2011;3:95–100. <https://doi.org/10.5897/JEBR11.009>.
46. Osman GY, Mohamed AM, Kader AA, Mohamed AA, Kader AA, Mohamed AA. Biological studies on Biomphalaria alexandrina snails treated with Furcraea sellosa marginata plant (family : Agavaceae) and Bacillus thuringiensis kurstaki (Dipel-2x). J Appl Pharm Sci 2011;1:47–55.
47. Bakry FA, Mohamed RT. Impact of Euphorbia milii latex on infectivity of Schistosoma mansoni larval stages to their hosts. Journal of Evolutionary Biology Research 2011;3:101–7. <https://doi.org/10.5897/JEBR11.015>.
48. Hassan S, Abdel Rahman E, Abd El Monem AR. Molluscicidal activity of butanol fraction of Meryta denhamii flowers against Lymnaea natalensis and Biomphalaria alexandrina. Glob Vet 2010;4:15–21.
49. Bakry FA. Use of some plant extracts to control Biomphalaria alexandrina snails with emphasis on some biological effects. Pestic Biochem Physiol 2009;95:159–65. <https://doi.org/10.1016/j.pestbp.2009.08.007>.
50. Attia WY, El-Bolkiny YE, Al-Sharkawi IM, Mohamed SH. Efficacy of Mandarin (Citrus Reticulata) Peel Extract in the Control of Schistosoma Mansoni Larval Stage and Their Intermediate Hosts. The Egyptian Society of Experimental Biology 2009;5:247–54.
51. Abdelgaleil SAM, El-Aswad AF, Nakatani M. Molluscicidal and anti-feedant activities of diterpenes from Euphorbia paralias L. Pest Manag Sci 2002;58:479–82. <https://doi.org/10.1002/ps.487>.
52. Motawea S, El-Gilany A-H, Salama O. Preliminary study of the molluscicidal action of Myrrh. Journal of Environmental Science 2001;21:153–62.
53. Allam AF, el-Sayad MH, Khalil SS. Laboratory assessment of the molluscicidal activity of Commiphora molmol (Myrrh) on Biomphalaria alexandrina, Bulinus truncatus and Lymnaea cailliaudi. J Egypt Soc Parasitol 2001;31:683–90.
54. El-Sayed MM, El-Nahas HA. Molluscicidal Steroidal Saponins From Yucca Elephantipes. Zagazig Journal of Pharmacological Science 2001;10:41–8.
55. Allam AF. Evaluation of different means of control of snail intermediate host of Schistosoma mansoni. J Egypt Soc Parasitol 2000;30:441–50.
56. Masoud AM, Fawzy SM, Salama OM. Laboratory studies on the molluscicidal and cercaricidal activities of Commiphora molmol. Egypt J Aquat Biol Fish 2000;4:251–66.
57. El-Ghaban AG, Eldiasty JG, M. H. M. Kamel O. Evaluation of Rosmarinus species extracted by different solvents against mosquito larvae, Biomphalaria species and different pathogenic bacteria. J Adv Sci Res 2015;6:44–9.
58. Jangwan JS, Aquino RP, Mencherini T, Picerno P, Singh R. Chemical constituents of ethanol extract of leaves and molluscicidal activity of crude extracts from Vitex trifolia Linn . Nerva Polonica 2013;59:19–32. <https://doi.org/10.2478/hepo-2013-0021>.
59. Mtemeli FL, Walter I, Tinago T, Shoko R. An assessment of the molluscicidal potential of Cucurbita maxima seed

- extracts on *Biomphalaria pfeifferi* and *Bulinus globosus* snails. *All Life* 2021;14:244–55. <https://doi.org/10.1080/26895293.2021.1901788>.
60. Isa M. Laboratory Assessment of Molluscicidal and Cercaricidal Activities of *Balanites aegyptiaca* against Vectors of Schistosomiasis (*Biomphalaria pfeifferi*). *International Journal of Research and Review* 2021;8:479–86. <https://doi.org/10.52403/ijrr.20210558>.
 61. Albagouri AH, Elegami AA, Koko WS. Molluscicidal Activity of Some Sudanese Medicinal Plants Against Vector Snails of *Schistosoma mansoni*. *Nile Journal for Agricultural Sciences* 2021;6:36–47.
 62. Abdelmageed E, Bushara HO, Abdelgadir M. Evidence of biological activity of *Pulicaria crispa* on *Biomphalaria pfeifferi* host snails of *Schistosoma mansoni*. *Biosciences Biotechnology Research Communications* 2017;11:55–9. <https://doi.org/10.21786/bbrc/11.1/8>.
 63. Mandefro B, Mereta ST, Tariku Y, Ambelu A. Molluscicidal effect of *Achyranthes aspera* L. (Amaranthaceae) aqueous extract on adult snails of *Biomphalaria pfeifferi* and *Lymnaea natalensis*. *Infect Dis Poverty* 2017;6:1–5. <https://doi.org/10.1186/s40249-017-0349-4>.
 64. Elsareh F, Abdalla R, Abdalla E. The effect of aqueous leaves extract of *Solenostemma argel* (Del Hayne) on egg masses and neonates of *Biomphalaria pfeifferi* snails. *Journal of Medicinal Plants* 2016;4:271–4.
 65. Mwonga KB, Waniki NE, Dorcas YS, Piero NM. Molluscicidal Effects of Aqueous Extracts of Selected Medicinal Plants from Makueni County, Kenya. *Pharm Anal Acta* 2015;6:1–4. <https://doi.org/10.4172/2153-2435.1000445>.
 66. Ahmed EA, Babiker OF, Abdalla RM. Molluscicidal Activity of Aqueous Leave Extract of *Solenostemma argel* (Del Hayne) on *Biomphalaria pfeifferi* Snails I : INTRODUCTION. *Journal of Basic and Applied Scientific Research* 2014;4:179–84.
 67. Hena JS, Sindama A, Kulawe D. Gas Chromatography Mass Spectrometric Analysis and Molluscicidal Effect of Crude Saponin from *Balanites aegyptiaca* (L .) Del . J Adv Biol Biotechnol 2017;15:1–7. <https://doi.org/10.9734/JABB/2017/28802>.
 68. Denou A, Togola A, Innjerdingen KT, Diallo D, Paulsen BS. Laboratory Assessment of Molluscicidal Activity of *Glinus Oppositifolius* (L .) Aug . DC . Related papers Laboratory Assessment of Molluscicidal Activity of. *International Journal of New Technology and Research* 2016;2:123–6.
 69. Kiros G, Erko B, Giday M, Mekonnen Y. Laboratory assessment of molluscicidal and cercariacidal effects of *Glinus lotoides* fruits. *BMC Res Notes* 2014;7:1–7. <https://doi.org/10.1186/1756-0500-7-220>.
 70. Molla E, Giday M, Erko B. Laboratory assessment of the molluscicidal and cercariacidal activities of *Balanites aegyptiaca*. *Asian Pac J Trop Biomed* 2013;3:657–62. [https://doi.org/10.1016/S2221-1691\(13\)60132-X](https://doi.org/10.1016/S2221-1691(13)60132-X).
 71. Michael ES, Yole D, Musila FM, Kutima H, Kareru P. Assessment of Molluscicidal , Cercaricidal and Miracidal Activities of Crude Extracts of *Azadirachta indica* and *Entada leptostachya*. *J Biol Agric Healthc* 2013;3:11–8.
 72. Otariho B, Morenikeji OA. Efficacy of aqueous and ethanolic extracts of leaves of *Chromolaena odorata* as molluscicide against different developmental stages of *Biomphalaria pfeifferi*. *Afr J Biotechnol* 2013;12:438–44. <https://doi.org/10.5897/AJB12.2156>.
 73. Oledibe PA, Morenikeji OA, Benson O. Zoology and Ecology Molluscicidal potency of *Ficus exasperata* (Vahl) against juvenile and adult *Biomphalaria pfeifferi*. *Zoology and Ecology* 2013;23:147–56. <https://doi.org/10.1080/21658005.2013.797150>.
 74. Benson O, Morenikeji OA. Molluscicidal Effects of Aqueous and Ethanolic Extracts of Lemongrass (*Cymbopogon Citratus*) Leaf against the Different Developmental Stages of *Biomphalaria Pfeifferi*. *New York Science Journal* 2012;5:70–7.
 75. Benson O. Efficacy of *Chromolaena odorata* leaf extracts on *Biomphalaria pfeifferi* eggs in control of Schistosomiasis. *Zoology and Ecology* 2012;22:236–9. <https://doi.org/10.1080/21658005.2012.745269>.
 76. Ismail MA, Koko WS, Osman EE, Dahab MM, Garbi MI, Alsadeg AM, et al. Molluscicidal Activity of *Acacia seyal* (Dell) Bark Methanolic Extract Against *Biomphalaria pfeifferi* Snails. *International Biological and Biomedical Sciences Journal* 2016;2:73–9.
 77. Ukwandu N, Bdaibo A, Okorie T, Nmorsi O. Molluscicidal effect of *Piper guineense*. *African Journal of Complementry Alternative Medicine* 2011;8:447–51. <https://doi.org/10.4314/ajtcam.v8i4.17>.
 78. Adenusi AA, Odaibo AB. A laboratory assessment of the potential molluscicidal activity of some Nigerian plant species used as anthelmintics. *Afr J Aquat Sci* 2010;35:251–8. <https://doi.org/10.2989/16085914.2010.545506>.
 79. Adentuji V o., Salawu OT. Efficacy of ethanolic leaf extracts of *Carica papaya* and *Terminalia catappa* as molluscicides against the snail intermediate hosts of schistosomiasis. *Journal of Medicinal Plants* 2010;4:2348–52. <https://doi.org/10.5897/JMPR10.468>.
 80. El-kamali HH, El-Nour OR, Khalid SA. Molluscicidal Activity of the Essential Oils of *Cymbopogon nervatus* Leaves and *Boswellia papyrifera* Resins. *Current Research Journal of Biological Sciences* 2010;2:139–42.
 81. Adenusi AA, Odaibo AB. Effects of varying concentrations of the crude aqueous and ethanolic extracts of *Dalbergia Sisso* plant parts on *Biomphalaria pfeifferi* egg masses. *African Journal of Traditional, Complementary and Alternative Medicines* 2009;6:139–49.
 82. Adenusi AA, Odaibo AB. Laboratory assessment of molluscicidal activity of crude aqueous and ethanolic extracts of *Dalbergia sissoo* plant parts against *Biomphalaria pfeifferi*. *Travel Med Infect Dis* 2008;6:219–27. <https://doi.org/10.1016/j.tmaid.2007.12.004>.
 83. Ojewole JAO, Nundkumar N, Adewunmi CO. MOLLUSCICIDAL, CERCARIACIDAL, LARVICIDAL AND ANTIPLASMODIAL PROPERTIES OF *BARRINGTONIA*

- RACEMOSA FRUIT AND SEED EXTRACTS. *Bol Latinoam Caribe Plantas Med Aromat* 2004;3:88–92.
84. Ahmed AAM, Ahmed SM. Extracts of Leaves and Seeds of the Neem Tree, *Azadirachta Indica*, As Environment-Oriented Molluscicides for Combating Schistosomiasis. *African Freshwater Malacology*, 2003, p. 235–49.
85. Diallo D, Marston A, Terreaux C, Toure Y, Smedsted Paulsen B, Hostettmann K. Screening of Malian Medicinal Plants for Antifungal, Larvicidal, Molluscicidal, Antioxidant and Radical Scavenging Activities. *Phytotherapy Research* 2001;15:401–6. <https://doi.org/10.1080/14786419.2014.931393>.
86. Wanyonyi AW, Chhabra SC, Mkoji G, Njue W, Tarus PK. Molluscicidal and antimicrobial activity of *Solanum aculeastrum*. vol. 74. 2003. [https://doi.org/10.1016/S0367-326X\(03\)00030-3](https://doi.org/10.1016/S0367-326X(03)00030-3).
87. Al-Zanbagi NA, Barrett J, Banaja AAA. Laboratory evaluation of the molluscicidal properties of some Saudi Arabian euphorbiales against *Biomphalaria pfeifferi*. *Acta Trop* 2001;78:23–9. [https://doi.org/10.1016/S0001-706X\(00\)00166-2](https://doi.org/10.1016/S0001-706X(00)00166-2).
88. Adewoyin FB, Adewunmi CO, Omisore NO, Olorunmola FO, Elusiyan CA, Agvedahunsi JM. Larvicidal, molluscicidal and antitrichomonal effects of methanolic extract of *Lycopersicon esculentum* mill leaf from south-west Nigeria. *Nigerian Journal of Natural Products and Medicines* 2012;16:29–32.
89. Agboola OI, Ajayi GO, Adesegun SA, Adesanya SA, Leavesò PÒ, Stemò A, et al. Comparative molluscicidal activities of fruit pericarp, leaves, seed and stem bark of *Blighia Unijugata* baker. *Pharmacognosy Journal* 2011;3:63–6. <https://doi.org/10.5530/pj.2011.25.11>.
90. Mata RCS, de Mendonca DIMD, Vieira L, Dos Santos AF, da Silva LA, Gaspar JF, et al. Molluscicidal Activity of Compounds Isolated from *Euphorbia conspicua* N. E. Br. *Journal of Brazilian Chemical Society* 2011;22:1880–7.
91. Changbunjong T, Wongwit W, Leemingsawat S, Tongtokit Y, Deesin V. Effect of crude extract of *Solanum xanthocarpum* against snails and mosquito larvae. *Southeast Asian Journal of Tropical Medicine and Public Health* 2010;41:320–5.
92. Baloch IB, Baloch MK, Baloch AK. Schistosomiasis suppressing deoxyphorbol esters from euphorbia cauducifolia L. Latex. *Planta Med* 2010;76:809–14. <https://doi.org/10.1055/s-0029-1240691>.
93. Medina JM, Peixoto JLB, Silva AA, Haraguchi SK, Falavigna DLM, Zamuner MLM, et al. Evaluation of the molluscicidal and *Schistosoma mansoni* cercariae activity of *Croton floribundus* extracts and kaurenoic acid. *Revista Brasileira de Farmacognosia* 2009;19:207–11. <https://doi.org/10.1590/S0102-695X2009000200005>.
94. Abouzid S, Orihara Y, Kawanaka M. Molluscicidal activity of polyacetylenes from *Ambrosia maritima* hairy roots. *Nat Prod Commun* 2007;2:177–80. <https://doi.org/10.1177/1934578x0700200214>.
95. Batista JJ, de Araújo HDA, Aguiar TW de A, Ferreira SA de O, Lima M de V, Pereira DR, et al. Toxic, cytotoxic and genotoxic effect of saline extract and fraction of *Parkia pendula* seeds in the developmental stages of *Biomphalaria glabrata* (Say 1818 – intermediate host) and cercaricide activity against the infectious agent of schistosomiasis. *Acta Trop* 2022;228. <https://doi.org/10.1016/j.actatropica.2022.106312>.
96. Pereira LPLA, Ribeiro ECG, Brito MCA, Araruna FOS, Araruna FB, Leite JAC, et al. Molluscicidal and cercaricidal activities of the essential oil of *Dysphania ambrosioides* (L.) Mosyakin & Clemants: Implications for the control of schistosomiasis. *Acta Trop* 2022;230. <https://doi.org/10.1016/j.actatropica.2022.106393>.
97. Silva YRR, Silva LD, Rocha TL, Dos Santos DB, Bezerra JCB, Machado KB, et al. Molluscicidal activity of *Persea americana* mill. (lauraceae) stem bark ethanolic extract against the snail *biomphalaria glabrata* (say, 1818): A novel plant-derived molluscicide? *An Acad Bras Cienc* 2020;92:1–16. <https://doi.org/10.1590/0001-37652020200715>.
98. Barros Gomes PR, Batista Reis J, Fernandes RP, Mouchrek Filho VE, De Souza AG, Fontenele MA, et al. Toxicity and molluscicidal activity of the essential oil *Pimenta dioica* against the snail *Biomphalaria glabrata*. *Rev Peru Biol* 2019;26:102–8.
99. Barros GRP, Oliveira MB, Sousa DA De, Caetano J, Fernandes RP, Louzeiro HC, et al. Larvicidal activity, molluscicide and toxicity of the essential oil of *Citrus limon* peels against, respectively, *Aedes aegypti*, *Biomphalaria glabrata* and *Artemia salina*. *Ecletica Quimica Journal* 2019;44:85–95. <https://doi.org/10.26850/1678-4618eqj.v44.4.p85-95>.
100. Faria RX, Rocha LM, Souza EPBSS, Almeida FB, Fernandes CP, Santos JAA. Molluscicidal activity of *Manilkara subsericea* (Mart.) dubard on *Biomphalaria glabrata* (Say, 1818). *Acta Trop* 2018;178:163–8. <https://doi.org/10.1016/j.actatropica.2017.11.012>.
101. Pereira LPLA, Dias CN, Miranda MiV, Firmo W da CA, Rosa C dos S, Brito MCA, et al. Molluscicidal effect of *Euphorbia umbellata* (Pax) Bruyns latex on *Biomphalaria glabrata*, *Schistosoma mansoni* host snail. *Journal of the Sao Paulo Institute of Tropical Medicine* 2017;40:1290–3. <https://doi.org/10.15537/smj.2019.12.24643>.
102. Sá JLF, Siqueira WN, Silva HAMF, Santos ML de O, Santos FTJ dos, Silva LRS, et al. Evaluation of molluscicidal activity of *Anadenanthera colubrina* extracts on adult mollusc and embryos of the species *Biomphalaria glabrata* (Say, 1818). *Scientia Plena* 2016;12:2–10. <https://doi.org/10.14808/sci.plena.2016.101001>.
103. [103] Rocha-filho CAA, Albuquerque LP, Silva LRS, Silva PCB, Coelho LCBB, Navarro DMAF, et al. Chemosphere Assessment of toxicity of *Moringa oleifera* flower extract to *Biomphalaria glabrata*, *Schistosoma mansoni* and *Artemia salina*. *Chemosphere* 2015;132:188–92. <https://doi.org/10.1016/j.chemosphere.2015.03.041>.
104. Feitosa A, Fonseca SA, César FA. A pentasubstituted pyridine alkaloid from the rhizome of *Jatropha elliptica* (Pohl) Muell. Arg. is active against

- Schistosoma mansoni and Biomphalaria glabrata. Parasitol Res 2014;113:1077–84. <https://doi.org/10.1007/s00436-013-3743-2>.
105. Santos CCS, Araújo SS, Santos ALLM, Almeida ECV, Dias AS, Damascena NP, et al. Evaluation of the toxicity and molluscicidal and larvicidal activities of schinopsis brasiliensis stem bark extract and its fractions. Revista Brasileira de Farmacognosia 2014;24:298–303. <https://doi.org/10.1016/j.bjp.2014.07.006>.
 106. Pereira Filho AA, França CRC, Oliveira D da S, Mendes RJ de A, Gonçalves J de RS, Rosa IG. Evaluation of the molluscicidal potential of hydroalcoholic extracts of Jatropha gossypifolia Linnaeus, 1753 on Biomphalaria glabrata. Rev Inst Med Trop Sao Paulo 2014;56:505–10. <https://doi.org/10.1590/S0036-46652014000600009>.
 107. Coelho da Silva CLPA, Vargas TS, Baptista DF. Molluscicidal activity of Moringa oleifera on Biomphalaria glabrata: Integrated dynamics to the control of the snail host of Schistosoma mansoni. Revista Brasileira de Farmacognosia 2013;23:848–50. <https://doi.org/10.1590/S0102-695X2013000500019>.
 108. Orechio P, Victor DM, Rapado LN, Sa A De, Fokoue HH, Scotti MT, et al. Schistosomiasis Control Using Piplartine against Biomphalaria glabrata at Different Developmental Stages. Neglected Tropical Diseases 2013;7:1–8. <https://doi.org/10.1371/journal.pntd.0002251>.
 109. Rodrigues KADF, Dias CN, Do Amaral FMM, Moraes DFC, Mouchrek Filho VE, Andrade EHA, et al. Molluscicidal and larvicidal activities and essential oil composition of Cymbopogon winterianus. Pharm Biol 2013;51:1293–7. <https://doi.org/10.3109/13880209.2013.789536>.
 110. Fontes-jr UR, Ramos CS, Serafini MR, Cavalcanti SCH, Alves PB, Lima GM, et al. Evaluation of the lethality of Porophyllum ruderale essential oil against Biomphalaria glabrata, Aedes aegypti and Artemia salina. African Journal of Biological Science 2012;11:3169–72. <https://doi.org/10.5897/AJB11.1314>.
 111. Lima MG, Augusto RDC, Vasconcellos MC De. Metabolic changes in Biomphalaria glabrata infected with Schistosoma mansoni exposed to latex of Euphorbia milii solution versus times of preparation. J Nat Prod 2012;5:222–32.
 112. Oliveira-filho EC, Geraldino BR, Coelho DR, Decarvalho RR, Paumgarten FJR. Comparative toxicity of Euphorbia milii latex and synthetic molluscicides to Biomphalaria glabrata embryos. Chemosphere 2010;81:218–27. <https://doi.org/10.1016/j.chemosphere.2010.06.038>.
 113. Teles T V., Bonfim RR, Alves PB, Blank AF, Jesus HCR, Quintans-Jr LJ, et al. Composition and evaluation of the lethality of Lippia gracilis essential oil to adults of Biomphalaria glabrata and larvae of Artemia salina. Afr J Biotechnol 2010;9:8800–4. <https://doi.org/10.5897/AJB10.113>.
 114. Santos NC, Dias CN, Coutinho-Moraes DF, Vilanova CM, Goncalves JDR, Souza N da S, et al. Toxicity and evaluation of molluscicidal activity of leaves of Turnera ulmifolia L. Brazilian Journal of Bioscience 2010;8:324–9.
 115. Silva TMS, Camara CA, Freire KRL, Da Silva TG, Agra MDF, Bhattacharyya J. Steroidal glycoalkaloids and molluscicidal activity of Solanum asperum Rich. fruits. J Braz Chem Soc 2008;19:1048–52. <https://doi.org/10.1590/S0103-50532008000500033>.
 116. Coutinho DF, Dias CS, Barbosa-Filho JM, Agra MF, Martins RM, Silva TMS, et al. Composition and molluscicidal activity of the essential oil from the stem bark of ocotea bracteosa (meisn.) mez. Journal of Essential Oil Research 2007;19:482–4. <https://doi.org/10.1080/10412905.2007.9699958>.
 117. Mello-Silva CC, de Vasconcellos MC, Pinheiro J, de Azevedo Rodrigues M de L. Physiological Changes in Biomphalaria glabrata Say, 1818 (Pulmonata: Planorbidae) caused by sub-lethal concentrations on the latex of Euphorbia splendens var. hislopiae N.E.B (Euphorbiaceae). Mem Inst Oswaldo Cruz 2006;101:3–8. https://doi.org/10.5005/jp/books/12167_1.
 118. Silva TMS, Câmara CA, Agra M de F, de Carvalho MG, Frana MT, Brandoline SVPB, et al. Molluscicidal activity of Solanum species of the Northeast of Brazil on Biomphalaria glabrata. vol. 77. 2006. <https://doi.org/10.1016/j.fitote.2006.05.007>.
 119. Silva TMS, Batista MM, Camara CA, Agra MF. Molluscicidal activity of some Brazilian Solanum spp. (Solanaceae) against Biomphalaria glabrata. Ann Trop Med Parasitol 2005;99:419–25. <https://doi.org/10.1179/136485905X36208>.
 120. Bezerra JCB, Silva IA, Ferreira HD, Ferri PH, Santos SC. Molluscicidal activity against Biomphalaria glabrata of Brazilian Cerrado medicinal plants. Fitoterapia 2002;73:428–30. [https://doi.org/10.1016/S0367-326X\(02\)00121-1](https://doi.org/10.1016/S0367-326X(02)00121-1).
 121. Meléndez PA, Capriles VA. Molluscicidal activity of plants from Puerto Rico. Ann Trop Med Parasitol 2002;96:209–18. <https://doi.org/10.1179/000349802125000600>.
 122. Dos Santos AF, Sant'Ana AEG. Molluscicidal properties of some species of Annona. Phytomedicine 2001;8:115–20. <https://doi.org/10.1078/0944-7113-00008>.
 123. Schall VT, Vasconcellos MC, Rocha RS, Souza CP, Mendes NM. The control of the schistosome-transmitting snail Biomphalaria glabrata by the plant Molluscicide Euphorbia splendens var. hislopiae (syn. milli Des. Moul.): a longitudinal field study in an endemic area in Brazil. Acta Trop 2001;79:165–70.
 124. Al-obaidi MJ, Abbas AH, Al-sa AJR, Al-husseiny IA. Schistosomiasis vector control using Cucumis melo Plant extracts with bioassay experiment 2018;18:425–30.
 125. Al-abidi MJ, Abbas AH, Khanon AY, Ibrahim K. Using the Thymus vulgaris extracts to control the snail vector

- in Schistosomiasis (Part II). The Iraqi Journal of Agricultural Sciences 2018;49:111–23.
126. Al-Obaidi MJL, Abbasp AH, Al-Azzaup AAM, Waaelp A. Survival rates of *Bulinus truncatus* as a way to determine the molluscicidal activity of *Ricinus Communis* extracts. Iraqi Journal of Biotechnology 2015;14:32–53.
127. El Babili F, Fabre N, Moulis C, Fouraste I. Molluscicidal activity against *Bulinus truncatus* of *Croton campestris*. vol. 77. 2006. <https://doi.org/10.1016/j.fitote.2006.03.003>.
128. Hannedouche S, Souchard JP, Jacquemond-Collet I, Moulis C. Molluscicidal and radical scavenging activity of quinones from the root bark of *Caryopteris x clandonensis*. vol. 73. 2002. [https://doi.org/10.1016/S0367-326X\(02\)00163-6](https://doi.org/10.1016/S0367-326X(02)00163-6).
129. Patel A V, Wright D, Blunden G, Sumner S, Rice J. Stable Molluscicide Formulation of an Aqueous Extract of *Euphorbia myrsinites*. Phytotherapy Research 2011;25:1412–4.
130. Apers S, Baronikova S, Sindambiwe JB, Witvrouw M, De Clercq E, Vanden Berghe D, et al. Antiviral, haemolytic and molluscicidal activities of triterpenoid saponins from *Maesa lanceolata*: Establishment of structure-activity relationships. Planta Med 2001;67:528–32. <https://doi.org/10.1055/s-2001-16489>.
131. Rug M, Ruppel A. Toxic activities of the plant *Jatropha curcas* against intermediate snail hosts and larvae of schistosomes. Tropical Medicine and International Health 2000;5:423–30. <https://doi.org/10.1046/j.1365-3156.2000.00573.x>.
132. Aminou HA, Mossalem HS, Alam-eldin Y. *Azadirachta indica* extracts on *Bulinus truncatus* snail and aquatic stages of *Schistosoma haematobium*. Parasitologists United Journal 2020:151–7. <https://doi.org/10.21608/puj.2020.39631.1085>.
133. Bakry FA, El-Garhy M, Abd El-Atti M, Th Atwa M. Effects of the Extracts of *Euphorbia Pulcherima* and *Atriplex Nummularia* on the Infectivity of *Schistosoma Haematobium* to *Bulinus Truncatus* Snails. Advances in Biomechanics 2017;1:34–41. <https://doi.org/10.11648/j.abm.20170102.12>.
134. Hasheesh WS, Mohamed RT, El-Monem SA. Biological and physiological parameters of *Bulinus truncatus* snails exposed to methanol extract of the plant *Sesbania sesban* plant. Adv Biol Chem 2011;01:65–73. <https://doi.org/10.4236/abc.2011.13009>.
135. Okeke, O.C. and Ubachukwu PO. Molluscicidal Effects of *Talinum triangulare* on *Bulinus truncatus*. Niger J Biotechnol 2011;22:13–6.
136. Bakry FA, Mohamed RT, Hasheesh WS. Impact of methanol extract of *Adenium obesum* plant on some biochemical and biological parameters of *Bulinus truncatus* snails. J Evol Biol Res 2011;3:87–94.
137. Larhsini M, Sebbane R, Kchakech H, Markouk M, Bekkouche K, Abbad A, et al. Screening of some Moroccan plant extracts for molluscicidal activity. Asian Journal of Experimental and Biological Sciences 2010;1:964–7.
138. Lahlou M, Berrada R. Potential of essential oils in schistosomiasis control in Morocco. International Journal of Aromatherapy 2001;11:87–96. [https://doi.org/10.1016/S0962-4562\(01\)80023-0](https://doi.org/10.1016/S0962-4562(01)80023-0).
139. Hmamouchi M, Lahlou M, Agoumi A. Molluscicidal activity of some Moroccan medicinal plants. Fitoterapia 2000;71:308–14. [https://doi.org/10.1016/S0367-326X\(99\)00152-5](https://doi.org/10.1016/S0367-326X(99)00152-5).
140. Suleiman J, Singh K, Bala AY, Muhammad MT, Abdullahi A. Comparison for the Effectiveness of Column Purified Fractions of *Allium cepa* Bulbs and *Allium sativum* Cloves against *Bulinus globosus* (Intermediate Host of Comparison for the Effectiveness of Column Purified Fractions of *Allium cepa* Bulbs and *Allium sativum*). Asian Journal of Advanced Research and Reports 2018;2:1–6. <https://doi.org/10.9734/ajarr/2018/v2i429774>.
141. Angaye TCN, Bassey SE, Ohimain EI, Izah SC, Asaigbe PI. Molluscicidal and synergicidal activities of the leaves of four Niger Delta Mangrove plants against schistosomiasis vectors. Journal of Environmental Treatment Techniques 2015;3:35–40.
142. Angaye TCN, Ohimain EI, Zige D V, Didi B, Biobelemoye N. Biocidal activities of Solvent Extracts of *Azadirachta indica* against Some Endemic Tropical Vector-borne Diseases. International Journal of Tropical Disease 2014;4:1198–208. <https://doi.org/10.9734/IJTDH/2014/11746>.
143. Okete J. Efficacy of the ethanolic extracts of *Talinum triangulare* for control of the fresh water snail, *Bulinus globosus*, the vector of urinary schistosomiasis. International Journal of Pure and Applied Zoology 2015;3:76–86.
144. Daniel UN, Onyemekara NN, Ohalet UN, Nnoli MC. Schistosomiasis snail-intermediate host control, using a natural plant. World J Pharm Sci 2013;2:52–63.
145. Salawu OT, Odaibo AB. The molluscicidal effects of *Hyptis suaveolens* on different stages of *Bulinus globosus* in the laboratory. Afr J Biotechnol 2011;10:10241–7. <https://doi.org/10.5897/AJB10.415>.
146. Olofintoye LK. Comparative evaluation of Molluscicidal effects of *Securidaca longepedunculata* (Fres.) and *Tephrosia bracteolata* (Guilland Perr) on *Bulinus globosus*. Journal of Parasitology and Vector Biology 2010;2:37–41. <https://doi.org/10.21608/eajbsf.2010.17449>.
147. Jia T-W, Wang W, Sun L-P, Lv S, Kun Y, Zhang N-M, et al. Molluscicidal effectiveness of Luo-Wei, a novel plant-derived molluscicide, against *Oncomelania hupensis*, *biomphalaria alexandrina* and *Bulinus truncatus*. Infect Dis Poverty 2019;27:70–9.
148. Ke W, Tu C, Cao D, Lin X, Sun Q, Zhang Q. Molluscicidal activity and physiological toxicity of quaternary benzo [c] phenanthridine alkaloids (QBAs) from *Macleaya cordata* fruits on *Oncomelania hupensis*. Neglected Tropical Diseases 2019:1–16.

149. Ludevese FL, Ludevese-Pascual G, Portugaliza HP. Molluscicidal activity of *Jatropha curcas* L. crude leaf extracts against *Oncomelania hupensis* quadrasi Mollendorf. *J Sci Eng Technol* 2018;186:177–86.
150. Ke W, Lin X, Yu Z, Sun Q, Zhang Q. Molluscicidal activity and physiological toxicity of *Macleaya cordata* alkaloids components on snail *Oncomelania hupensis*. *Pestic Biochem Physiol* 2017;143:111–5. <https://doi.org/10.1016/j.pestbp.2017.08.016>.
151. Han BX, Chen J. Acacetin-7-rutinoside from *buddleja lindleyana*, A new molluscicidal agent against *oncomelania hupensis*. *Zeitschrift Fur Naturforschung - Section C Journal of Biosciences* 2014;69 C:186–90. <https://doi.org/10.5560/ZNC.2013-0179>.
152. Yang F, Long E, Wen J, Cao L, Zhu C, Hu H, et al. Linalool, derived from *Cinnamomum camphora* (L.) Presl leaf extracts, possesses molluscicidal activity against *Oncomelania hupensis* and inhibits infection of *Schistosoma japonicum*. *Parasit Vectors* 2014;7:1–13.
153. Li X, Shan X, Pan J, Yu P, Mao Z. Effects of the molluscicidal agent GA-C 13: 0, a natural occurring ginkgolic acid, on snail mitochondria. *Pestic Biochem Physiol* 2012;103:115–20. <https://doi.org/10.1016/j.pestbp.2012.04.006>.
154. Chen YQ, Xu QM, Liu YL, Li XR, Yang SL, Zhuge HX. Laboratory evaluation of the molluscicidal activity of *pulsatilla Chinensis* (Bunge) regel saponins against the snail *oncomelania hupensis*. *Biomedical and Environmental Sciences* 2012;25:224–9. <https://doi.org/10.3967/0895-3988.2012.02.015>.
155. Sun Q, Yang Y, Zhang Q. Effect and mechanism of *Sapium sebiferum* against snail (*Oncomelania hupensis*) and control of schistosomiasis. *International Conference on Biomedical Engineering and Biotechnology* 2012:190–3. <https://doi.org/10.1109/iCBEB.2012.165>.
156. [156] Ming Z, Gui-Yin L, Jian-Guo Z, Li Z, Ke-Long H, Jin-Ming S, et al. Evaluation of molluscicidal activities of benzo[c]phenanthridine alkaloids from *Macleaya cordata* (Willd) R. Br. on snail hosts of *Schistosoma japonicum*. *Journal of Medicinal Plants Research* 2011;5:521–6.
157. Peng F, Liu M, Huang Q, Liu N, Yang H, Sun H, et al. Molluscicidal Effect of *Eomecon chionantha* alkaloids against *Oncomelania hupensis* snails. *Southeast Asian Journal of Tropical Medicine and Public Health* 2011;42:289–96.
158. Li Y, Chen J, Han B, Pan G, Guo D. Molluscicidal activities of *Ginkgo biloba* leaf against the snail *Oncomelania hupensis*. *Journal of Medicinal Plants Research* 2010;4:2466–72. <https://doi.org/10.5897/jmpr10.185>.
159. Han BX, Chen J, Yang X, Wang S, Li CG, Han FA. Molluscicidal activities of medicinal plants from eastern China against *Oncomelania hupensis*, the intermediate host of *Schistosoma japonicum*. *Brazilian Journal of Pharmacognosy* 2010;20:712–7. <https://doi.org/10.1590/S0102-695X2010005000018>.
160. Zou FC, Duan G, Xie YJ, Zhou Y, Dong GD, Lin RQ, et al. Molluscicidal activity of the plant *Eupatorium adenophorum* against *Oncomelania hupensis*, the intermediate host snail of *Schistosoma japonicum*. *Ann Trop Med Parasitol* 2009;103:549–53. <https://doi.org/10.1179/136485909X451780>.
161. Yang X ming, Chen S xia, Xia L, Chen J. Molluscicidal activity against *Oncomelania hupensis* of *Ginkgo biloba*. *Fitoterapia* 2008;79:250–4. <https://doi.org/10.1016/j.fitote.2007.11.030>.
162. Ke W, Yang J, Meng Z, Ma A. Evaluation of molluscicidal activities of *Arisaema tubers* extracts on the snail *Oncomelania hupensis*. *Pestic Biochem Physiol* 2008;92:129–32. <https://doi.org/10.1016/j.pestbp.2008.07.003>.
163. Chen SX, Wu L, Yang XM, Jiang XG, Li LG, Zhang RX, et al. Comparative molluscicidal action of extract of *Ginkgo biloba* sarcotesta, arecoline and niclosamide on snail hosts of *Schistosoma japonicum*. *Pestic Biochem Physiol* 2007;89:237–41. <https://doi.org/10.1016/j.pestbp.2007.07.010>.
164. Wei FH, Xu XJ, Liu JB, Dai YH, Dussart G, Trigwell J. Toxicology of a potential molluscicide derived from the plant *Solanum xanthocarpum*: A preliminary study. *Ann Trop Med Parasitol* 2002;96:325–31. <https://doi.org/10.1179/000349802125000727>.
165. Rug M, Ruppel A. Toxic activities of the plant *Jatropha curcas* against intermediate snail hosts and larvae of schistosomes. *Tropical Medicine and International Health* 2000;5:423–30. <https://doi.org/10.1046/j.1365-3156.2000.00573.x>.
166. Sukumaran D, Parashar BD, Rao KM. Evaluation of some plant molluscicides against a freshwater snail *Lymnaea luteola*, the vector of animal schistosomiasis evaluation of some plant molluscicides against a freshwater snail *Lymnaea luteola*, the vector of animal schistosomiasis. *Pharm Biol* 2002;40:450–5. <https://doi.org/10.1076/phbi.40.6.450.8448>.
167. Samah MA-K, Samah NE, Ismail ShA. Molluscicidal effects of acetone and ethanol extracts of clove (*Syzygium aromaticum*) against *Monacha cartusiana* (Gastropoda: Hygromiidae) snails under laboratory and field conditions at Sharkia Governorate. *Egyptian Journal of Plant Protection Research Institute* 2020;3:595–603.
168. El-Atti MA, A.A E, A.M. K, Elgohary WS. Control of the glassy clover snails *Monacha cartusiana* using *Zingiber officinale* extract as an ecofriendly molluscicide Mahmoud. *African Journal of Biological Science* 2019;15:101–15.
169. Abdel-mogib M, Dawidar AEM, Mortada MM, Raghieb HM. against *Monacha cartusiana* molluscicidal activity of *Balanites aegyptiaca* against *Monacha cartusiana*. *Pharm Biol* 2012;50:1326–9. <https://doi.org/10.3109/13880209.2012.674950>.
170. Abdelhady Ahmed EA. EFFECT OF CASTOR PLANT (*RICINUS COMMUNIS*) EXTRACTS ON THE TERRESTRIAL SNAIL *MONACHA OBSTRUCTA* (*FERUSSAC*). *Plant Arch* 2020;20:1734–42.

171. Hussien Ali IH, Mohammed El-Sayed SA, Ali Fathey R, Abd El-Halim SM. Molluscicidal and antifeedant impacts of some weed extracts on the land snail *Monacha obstructa* (Pfeiffer). *Fayoum Journal of Agricultural Research and Development* 2020;34:204–17.
172. Alzabib AA, Abobakr Y, Al-sarar AS, Hussein HI, Basudan OA, El-gamal AA, et al. Molluscicidal activity of cardiac glycosides isolated from *Adenium obesum*. *Pest Manag Sci* 2019;75:2770–5. <https://doi.org/10.1002/ps.5388>.
173. Preet S. Laboratory evaluation of Molluscicidal and Cercaricidal Potential of *Artemisia annua* (Family: Asteraceae). *Ann Biol Res* 2010;2:373–83.
174. Muraleedharan K, Murthy SHK. Preliminary screening of leaves of *Leucas aspera* and *Parthenium hysterophorus* against fresh water snail, *Indoplanorbis exustus*. *Indian Journal of Veterinary Research* 2009;17:8–12.
175. Al-sarar A, Hussein H, Abobakr Y, Bayoumi A. Molluscicidal Activity of Methomyl and Cardenolide Extracts from *Calotropis procera* and *Adenium arabicum* Against the Land Snail *Monacha cantiana*. *Molecules* 2012;17:5310–8. <https://doi.org/10.3390/molecules17055310>.
176. Usman AM, Shinkafi SA. Molluscicidal Activity of Some Selected Plants on Freshwater Snail *Lanistes Ovum*. *International Journal of Science for Global Sustainability* 2016;2:24–37.
177. Ali TH, Abdul Jabbar Hamed A. The Molluscicidal Activity of *Melia azadirachta* on The Fresh water snail *Physa acuta* (Draparnaud , 1805) The Molluscicidal Activity of *Melia azadirachta* on The Fresh water snail *Physa acuta* (Draparnaud , 1805). *Journal of Pure and Applied Sciences* 2019;5:1–7.
178. Dib R, Makhoul K, Maalouf R. Preliminary bioactivity investigation of *Styrax officinalis* fruit extract as potential biopesticide. *Journal of Pharmacognosy and Phytotherapy* 2016;8:209–13. <https://doi.org/10.5897/JPP2016.0422>.
179. Souza BA De, Cristina L, Chicarino ED, Cristina E, Bessa DA. Phytochemical and Molluscicidal activity of *Mikania glomerata* Sprengel (Asteraceae) in Different Lifestages of *Subulina octona* (Mollusca , Subulinidade). *Brazilian Archives of Biology and Technology* 2014;57:261–8.
180. Souza BA, Silva LC Da, Chicarino ED, Bessa ECA. Preliminary phytochemical screening and molluscicidal activity of the aqueous extract of *Bidens pilosa* Linné (Asteraceae) in *Subulina octona* (Mollusca , Subulinidade). *Annals of the Brazilian Academy of Sciences* 2013;85:1557–66.
181. Akinpelu BA, A. DC, I. AF, O. IE, O.O. O. Effect of stem - bark of *Erythrophleum suaveolens* (Guill. Perri.) saponin on fresh water snail (*Lanistes lybicus*) tissues. *Afr J Environ Sci Tech* 2012;6:446–51. <https://doi.org/10.5897/ajest12.007>.
182. Silva L, Durço E, Pinheiro J, Bessa ECDA. Effect of *Solanum paniculatum* leaf extract on food consumption , fertility and carbohydrate metabolism of *Bradybaena similis* snail. *J Nat Prod* 2013;6:168–76.
183. Kariuki ST, Kariuki JM, Mailu BM, Muchiri DR. *Phytolacca octandra* (L.), *Phytolacca dodecandra* (LHerit) and *Balanites aegyptiaca* (L.) extracts as potential molluscicides of schistosomiasis transmitting snails. *Journal of Medicinal Plants Research* 2016;10:823–8. <https://doi.org/10.5897/jmpr2016.6237>.
184. Nnamdi UC, Chuka EP, Chukwudi UK, Chinelo U, Okwudiri IM. Molluscicidal Assessment of Aqueous Extract of *Moringa oleifera* Lam seed on *Bulinus* Snail for the control of Schistosomiasis Molluscicidal Assessment of Aqueous Extract of *Moringa oleifera* Lam Seed on *Bulinus* Snail for the Control of Schistosomiasis. *South Asian Journal of Research in Microbiology* 2020;7:31–9. <https://doi.org/10.9734/SAJRM/2020>.
185. Mtemeli FL, Walter I, Shoko R. Molluscicidal effects of pumpkin seed extracts on *Schistosoma* vectors 2020:1–13.
186. Eisa AI, Mohammed Elfaki TE, Abdellah MA, Elfaki M. Molluscicidal Activity of *Khaya Grandifoliola* Extracts Against Freshwater Snails in Khartoum State- Sudan. *International Journal of Academic Health and Medical Research* 2019;3:1–4.
187. Kindiki MG. Molluscicidal Activity of Selected Plant Extracts. 2014.
188. Caixeta SC, Magalhães LG, de Melo NI, L. Wakabayashi KA, Paula Aguiar G de, Paula Aguiar D de, et al. Chemical Composition and *in Vitro* Schistosomicidal Activity of the Essential Oil of *Plectranthus Neochilus* Grown in Southeast Brazil. *Chem Biodivers* 2011. <https://doi.org/10.1002/cbdv.201100167>.
189. Soares MH, Dias HJ, Vieira TM, Mendes Souza MG, Cruz AF, Badoco FR, et al. Chemical Composition, Antibacterial, Schistosomicidal, and Cytotoxic Activities of the Essential Oil of *Dysphania Ambrosioides* (L.) *Mosyakin* & *Clematis* (Chenopodiaceae). *Chem Biodivers* 2017. <https://doi.org/10.1002/cbdv.201700149>.
190. Domínguez-Vigil IG, Mata-Cárdenas BD, Esquivel-Ferriño PC, Ávalos-Alanís FG, Vargas-Villarreal J, Rayo Camacho-Corona M del. Antigiardial Activity of *Foeniculum Vulgare* Hexane Extract and Some of Its Constituents. *Plants* 2022. <https://doi.org/10.3390/plants11172212>.
191. Tripathi SM, Singh DK. Molluscicidal activity of *Punica granatum* bark and *Canna indica* root. *Brazilian Journal of Medical and Biological Research* 2000;33:1351–5. <https://doi.org/10.1590/S0100-879X2000001100014>.
192. Dossaji S, Kubo I. Quercetin 3-(2"-Galloyl)glucoside), A Molluscicidal Flavanoid from *Polygonum Senegalense*. *Phytochemistry* 1980;19:482.
193. Marston A, Hostettmann K. Plant molluscicides. *Phytochemistry* 1985;24:639–52. [https://doi.org/10.1016/S0031-9422\(00\)84870-0](https://doi.org/10.1016/S0031-9422(00)84870-0).

194. Doctor TR. Molluscicidal Performance of Kae Farang (*Gliciridia sepium*, JacqStrudel and Siam Weed (*Chromolaena odorata*, Linn.) Leaf Extracts on Golden Snail (*Pomacea caniculata*, Linn.) 2019.
195. Singh V, Kumar P, Singh VK, Singh DK. Effect of abiotic factors on the molluscicidal activity of oleoresin of *Zingiber officinale* against the snail *Lymnaea acuminata*. *Nat Sci (Irvine)* 2010;02:1148–54. <https://doi.org/10.4236/ns.2010.210142>.
196. Abubakar A, Bala AY, Singh K. Plant Molluscicides and their Modes of Action: a Review. *International Journal of Advanced Scientific Research* 2017;2:37–49.
197. Sharifi-Rad J, Sureda A, Tenore GC, Daglia M, Sharifi-Rad M, Valussi M, et al. Biological Activities of Essential Oils: From Plant Chemoecology to Traditional Healing Systems. *Molecules* 2017. <https://doi.org/10.3390/molecules22010070>.
198. Kalinda C, Chimbari MJ, Grant WE, Wang H, Odhiambo JN, Mukaratirwa S. Simulation of Population Dynamics of *Bulinus Globosus*: Effects of Environmental Temperature on Production of *Schistosoma Haematobium Cercariae*. *PLoS Negl Trop Dis* 2018. <https://doi.org/10.1371/journal.pntd.0006651>.
199. Henderson I, Triebskorn R. Chemical control of terrestrial gastropods. *Molluscs as Crop Pests* 2009:1–31. <https://doi.org/10.1079/9780851993201.0001>.
200. Hamed M, El-Amin S, Abdel-Fattah AS. Effect of some environmental conditions on the molluscicidal activity of *yucca filamentosa* “marginata” and *cestrum purpureum*. *Pharmacologyonline* 2015;2:142–7.